

Ecosystem health in Wellington City urban streams:

Stage one summary report

E Harrison
Environmental Science Department

For more information, contact the Greater Wellington Regional Council:

Wellington
PO Box 11646

T 04 384 5708
F 04 385 6960
www.gw.govt.nz




Masterton
PO Box 41

T 06 378 2484
F 06 378 2146
www.gw.govt.nz

GW/ESCI-T-19/123

November 2019

www.gw.govt.nz
info@gw.govt.nz

Report prepared by:	E Harrison	Senior Freshwater Scientist	
Report reviewed by:	S Greenfield	Freshwater Scientist, Ripple Ecology	
Report approved for release by:	M Oliver	Team Leader, Marine and Freshwater	 Date: 19/11/19

DISCLAIMER

This report has been prepared by Environmental Science staff of Greater Wellington Regional Council (GWRC) and as such does not constitute Council policy.

In preparing this report, the authors have used the best currently available data and have exercised all reasonable skill and care in presenting and interpreting these data. Nevertheless, GWRC does not accept any liability, whether direct, indirect, or consequential, arising out of the provision of the data and associated information within this report. Furthermore, as GWRC endeavours to continuously improve data quality, amendments to data included in, or used in the preparation of, this report may occur without notice at any time.

GWRC requests that if excerpts or inferences are drawn from this report for further use, due care should be taken to ensure the appropriate context is preserved and is accurately reflected and referenced in subsequent written or verbal communications. Any use of the data and information enclosed in this report, for example, by inclusion in a subsequent report or media release, should be accompanied by an acknowledgement of the source.

The report may be cited as:

Harrison, E. 2019. Ecosystem health in Wellington City urban streams. Greater Wellington Regional Council, Publication No. GW/ESCI-T-19/123, Wellington.

Executive summary

There is an extensive network of streams in the Wellington City area. Many of these streams have been piped and historically there has been minimal information on their ecological health (in terms of habitat condition, macroinvertebrate community health and fish populations). To address this knowledge gap, Greater Wellington Regional Council (GWRC) has worked with Wellington City Council (WCC) over the last three years to better understand the ecological health of Wellington's piped and open urban streams.

Monitoring commenced in 2016 and has continued over the summer months until 2019. During this time, a range of open urban streams in Wellington City have been surveyed on a single occasion to assess habitat condition, macroinvertebrate community health and fish populations. During this period habitat condition, macroinvertebrate and fish populations were also surveyed in piped streams by EOS Ecology.

The purpose of the monitoring is to provide a baseline of information from which a long-term monitoring programme for Wellington City stream ecosystem health can be developed. This monitoring programme is needed to report against National Policy Statement for Freshwater Management and GWRC Proposed Natural Resources Plan objectives for ecosystem health. Information is also needed to inform a range of Wellington City Council and Wellington Water strategies aimed at reducing the impact of urban land use on Wellington City stream ecosystems. This report presents the results of baseline monitoring for open streams. Results for the piped stream survey are attached in separate reports.

Ecosystem health data collected to date indicate Wellington City urban streams contain a range of values related to habitat, macroinvertebrates and fish. Key findings were:

- Macroinvertebrate communities were generally in better condition in streams with less urban landuse and impervious area in the upstream catchment.
- Fish communities within urban streams were generally in fair to poor condition. However, there is greater fish diversity in sections of stream with direct connection to the sea.
- Inanga spawning was occurring in bank vegetation in the lower Kaiwharawhara Stream.
- Banded kokopu and eel species were the most abundant fish species present across the majority of sites surveyed. Native fish species present included at risk and declining species such as koaro, inanga, redfin bully, longfin eel and giant kokopu.
- Barriers to fish movement are likely to be one of the major pressures influencing fish communities in Wellington urban streams. The only fish species found upstream of piped sections of stream are climbing species (banded kokopu, koaro and eel species). The presence of banded kokopu and eels in piped stream sections was confirmed in the piped stream survey.

- Macroinvertebrate and fish communities in Wellington City streams appear to be driven by different environmental stressors. The healthiest macroinvertebrate communities were generally found higher in catchments where habitat and water quality degradation from urban runoff is less. In contrast, fish communities were generally in better condition at sites lower in the catchment where there were likely to be less barriers to migration to and from the sea. This finding highlights the need for a whole of catchment approach for improving ecosystem health of Wellington City streams.
- The initial piped stream survey has shown that some sections of piped streams have habitat which supports fish and macroinvertebrate life, however species richness and abundance is greater in free flowing streams. The complete enclosing of open channels significantly reduces habitat quality for stream life and means only a subset of available taxa within catchments are able to persist there.

The next stage will involve design of a long term monitoring programme of Wellington City stream ecosystem health. This will involve working with WCC to identify objectives of the programme, monitoring sites, parameters to be measured, sampling frequency and reporting required. Targeted investigations needed to answer specific questions around management actions will also be identified. The programme will be designed in 2019/20 for implementation in 2020/21.

Contents

Executive summary	i
1. Introduction	1
1.1 Background and objectives	1
1.2 Linkages to management of urban streams in Wellington City	2
2. Methods	4
2.1 Site location and data collection	4
2.2 Habitat Assessments	6
2.3 Macroinvertebrates	7
2.4 Fish	7
2.5 Data analysis	8
2.5.1 Impervious and vegetated area within each catchment	8
2.5.2 Macroinvertebrate and fish community structure	8
3. Results	9
3.1 Habitat condition	9
3.1.1 Macroinvertebrates	10
3.1.2 Fish and koura	15
4. Discussion and conclusions	22
4.1 Urban stream ecosystem health and implications for urban stream management in Wellington City	22
4.2 Next steps for Wellington urban stream monitoring	24
Acknowledgements	26
References	27
Appendix 1: Photos from fish sampling	28
Appendix 2: Freshwater Ecology of Piped Streams in Wellington: Pilot Study Interim and Final Reports	33

1. Introduction

1.1 Background and objectives

There is an extensive network of streams within the Wellington City area. Of the rivers and streams monitored in the Wellington region, those with significant urban landuse within their catchments are consistently identified as having the poorest water quality and ecological health (eg. Mitchell and Heath, 2019). Pressures on these urban streams include:

- Contamination and erosion from stormwater discharges
- Contamination from sewer overflows/cross connections and other unauthorised discharges
- Habitat degradation through straightening and lining of stream banks with concrete or rock armouring
- Loss of stream habitat and function through stream piping and reclamation
- Barriers to fish migrations such as weirs and perched culverts

Water and habitat quality in urban streams is frequently poor, reflecting the land use activities in surrounding urban catchments. This in turn affects biota, the ability of people to use waterways for recreation, and degrades downstream receiving environments. The poor ecological health of urban stream systems is commonly referred to as “urban stream syndrome” (Walsh et al., 2005), where several physical, chemical and biological characteristics of urban streams are altered. Urban stream health has been shown to be strongly correlated to the amount of impervious surfaces such as roofs, roads and carparks in the catchment (Walsh et al., 2005).

Previous analyses have shown that urban stream health in the Wellington region (indicated by macroinvertebrate communities) decreases to fair/poor condition with increasing impervious land cover greater than approximately 15% (Warr, 2009). However, to date, there is not a full understanding of the range of ecosystem health values related to habitat, fish and macroinvertebrates across the range of Wellington urban streams (James, 2015). Nor is there any ecological information on the habitat quality, macroinvertebrates and fish in the piped stream systems across Wellington City (James, 2015). This information is required to inform the decision making related to managing the effects of land development and stormwater management.

Greater Wellington Regional Council (GWRC) is implementing a programme to identify and monitor the ecosystem health values of Wellington City’s urban streams. The programme includes assessments in open and piped streams and is jointly funded by Wellington City Council (WCC) and GWRC.

Stage one monitoring commenced in 2016 and continued over the summer months until 2019. During this time, a range of open and piped streams in Wellington City have been surveyed to assess habitat condition, macroinvertebrate community health and fish populations. (see the piped streams project reports attached in Appendix 2). The intention of this monitoring was to provide a baseline assessment of ecosystem health, which

would then inform a long-term monitoring programme to measure the state and trends in the ecosystem health of urban streams in Wellington City (Stage two).

This monitoring will be used to report against the objectives in the National Polity Statement for Freshwater Management and Proposed Natural Resources Plan (PNRP) for the Greater Wellington Region. The monitoring results will be used to assess the effectiveness of Wellington City Council strategies which aim to decrease the impacts of urban land use and stormwater runoff on Wellington City streams.

This report presents the results of Stage one monitoring of WCC urban streams. The report also explores the likely effects of urbanisation on stream ecosystem health in Wellington City streams. Finally, next steps for long term monitoring of state and trends in the ecosystem health of streams in Wellington City (stage two) are presented.

1.2 Linkages to management of urban streams in Wellington City

There are several management strategies and policies which require information on the current state of ecosystem health of Wellington City urban streams and seek to reduce the effect of surrounding urban land use (Table 1.1).

Assessments of urban stream ecosystem health using habitat, macroinvertebrate and fish will support these policies and strategies by:

- Identifying the range of biodiversity values in Wellington City's urban streams related to fish and macroinvertebrate communities;
- Identifying habitat requirements for the management of healthy fish and macroinvertebrate communities in urban environments;
- Providing reliable data for community groups to target areas for restoration activities;
- Raising awareness of urban stream biodiversity among local communities
- Understanding the current distribution of fish species within the city and changes from historical distributions;
- Assessing the effectiveness of management actions; and
- Understanding how and where to prioritize actions such as improvements to fish passage in open and piped streams.

The monitoring programme will be designed to provide as much information as possible to inform these policies and strategies. However, it won't be possible to answer all questions about the effect of urban land use in WCC on stream ecosystem health. In some cases targeted studies may need to be set up to answer specific questions such as the effectiveness of stormwater action plans.

Table 1.1: Strategies and policies related to the management of urban stream ecosystem health in Wellington City

Agency	Strategy/Policy	Linkage
Greater Wellington Regional Council	Proposed Natural Resources Plan and National Policy Statement for Freshwater Management	Assessment of objectives related to ecosystem health
Wellington City Council	Our Natural Capital	Identification of biodiversity values and maintaining/improving these values into the future
Wellington City Council	Resilience Strategy	Understanding of current urban freshwater ecosystem health and effects of any actions to improve stormwater quality entering urban streams
Wellington City Council	Urban Growth Strategy	Understanding the effects of future urban development options or urban stream ecosystem health to minimise any impacts
Wellington City Council/New Zealand Transport Authority	Roading strategies	Effects of roading maintenance and construction on urban stream ecosystem health
Wellington Water/Wellington City Council	Stormwater management plans	Assessment of stormwater management actions related to the improvement of urban stream ecosystem health
Wellington Water	Stormwater system maintenance	Information to assist with the maintenance of stormwater assets and allowances for fish passage through piped streams

2. Methods

2.1 Site location and data collection

Fifty one survey sites were spread across a range of urban streams throughout the WCC area, including sites upstream of urban areas and stream sections within the current urban footprint. Sites were also located in catchments with no current urban land use (Waipapa Stream catchment on the Wellington South Coast; South Makara stream catchment) (Figure 2.1). Data collected at each site is outlined in Table 2.1.




Figure 2.1: Locations of Wellington City stream sites surveyed for habitat quality, macroinvertebrate and fish community health between 2016 and 2019 (see Table 2 for site names and details)

Table 2.1: Location of survey sites and data collected at each site (✓ = collected 2018/19; ✓ = collected 2017/18; ✓ = collected 2016/17)

Catchment	Site code	Site name	Habitat	Macro-invertebrates	Fish	
					Electro Fishing	Spotlight
Karori	K1	Karori S at Karori Pk	✓	✓	✓	✓
	K2	Karori S at Castlemaine Cl	✓	✓	✓	✓
	K4	Karori S at Makara Peak Mt Bike Pk	✓	✓	✓	
	K5	Karori S at Makara coast	✓	✓	✓	✓
	K6	South Makara S tributary at South Makara Rd	✓	✓	✓	✓
	K6a	South Makara S at South Makara Rd				✓
	K7	Karori S at South Karori Rd	✓	✓	✓	
	K8	Karori S opposite Sunshine Ave	✓	✓	✓	
	K9	Karori S at Darwin St	✓	✓	✓	✓
	K10	Karori S at Futuna Cl			✓	✓
Ngauranga	N1	Ngauranga S at Newlands Rd	✓	✓	✓	
	N2	Ngauranga S near Alex Moore Pk	✓	✓	✓	
	N3	Ngauranga S at Mount Kaukau	✓	✓	✓	
	N4	Ngauranga S at Tyers Stream Reserve	✓	✓	✓	
	N5	Ngauranga S at Taylor Prestons	✓	✓	✓	
	N6	Ngauranga S at Ngauranga Gorge rail crossing				✓
	N7	Ngauranga S 400m above mouth	✓	✓	✓	✓
Moturoa	M1	Moturoa S at bottom of Central Pk	✓	✓	✓	
	M2	Moturoa S at top of Central Pk	✓	✓	✓	
Kaiwharawhara	Kai1	Kaiwharawhara S below the dam	✓✓	✓	✓	✓
	Kai2	Kaiwharawhara S below piped section	✓✓	✓	✓	✓
	Kai3	Kaiwharawhara S at Otari Wilton's Bush	✓✓	✓	✓✓	✓
	Kai4	Kaiwharawhara S below Korimako confluence	✓	✓		✓
	Kai5	Korimako S at Girl Guides	✓✓	✓	✓	✓
	Kai6	Kaiwharawhara S at Ngaio Gorge	✓	✓		✓
	KaiSpt	Kaiwharawhara S at Spotlight				✓✓
Owhiro	O1	Owhiro S upstream of TNT Landfill	✓	✓	✓	✓
	O1a	Owhiro S on Ohiro Rd below Brooklyn town				✓
	O1b	Owhiro S headwaters at Elliot Pk				✓

Table 2.1 cont: Location of survey sites and data collected at each site (✓ = collected 2018/19; ✓ = collected 2017/18; ✓ = collected 2016/17)

Catchment	Site code	Site name	Habitat	Macro-invertebrates	Fish	
					Electro Fishing	Spotlight
Owhiro	O2a	Owhiro S downstream of TNT Landfill	✓	✓	✓	✓
	O2b	Owhiro S below Landfill Rd Confluence				✓
	O2c	Owhiro S at S landfill gate on Landfill Rd				✓
	O3	Owhiro S upstream of Murchison St	✓✓	✓	✓	✓
	O4a	Owhiro S downstream of Happy Valley Rd	✓	✓	✓	✓
Kumutoto	KM01	Kumutoto Stream at Vic Uni	✓	✓		✓
Lyaill Bay	LB01	Ngaroma Stream	✓	✓		✓
Mahanga	MH01	Mahanga Bay Stream	✓	✓		✓
Miramar	MR01	Maupuia Stream	✓	✓		✓
Paikawakawa	PA01	Island Bay Stream trib at Farnham St	✓	✓		✓
	PA02	Island Bay Stream trib at Mana Karioi	✓	✓		✓
Pipitea	PT01	Puketea Stream at Botanical Gardens	✓	✓	✓	✓
	PT02	Pipitea Stream at Botanical Gardens below pond			✓	
Te Poti	TP01	Te Poti Stream	✓	✓		✓
Waimapihi	WA01	Waimapihi Stream	✓	✓	✓	
	WA02	Clinical Track Stream	✓	✓		✓
	WA03	Polhill Stream	✓	✓		✓
Waipapa	WP01	Waipapa Upstream	✓	✓	✓	✓
	WP02	Waipapa Stream at coast	✓	✓	✓	
Waitangi	WT01	Bells Rd Stream	✓	✓		✓
	P1	Papawai S at Prince of Wales Pk	✓	✓	✓	
Wadestown	WD1	Wadestown Stream				✓

2.2 Habitat Assessments

Habitat assessments were completed at 41 sites (once at 36 sites, twice at 5 sites – Table 2.1) following methods outlined in Clapcott (2015). Habitat assessments were only collected at sites where macroinvertebrates were collected which covered the range of stream types and urban impacts across Wellington City. The assessment provides an indication of the physical stream habitat condition and its ability to support stream biota. The assessment incorporates the following ten variables: deposited sediment cover,

macroinvertebrate habitat abundance and diversity, fish habitat abundance and diversity, hydraulic heterogeneity, bank erosion and vegetation, and riparian width and shade. Each category is scored between 1 ('poor') and 10 ('excellent'). Summation of individual scores provides an overall total habitat quality score for each site (lowest and highest possible scores are 10 and 100, respectively).

2.3 Macroinvertebrates

Macroinvertebrates were sampled at 41 sites with hard bottom substrate on a single occasion (Table 2.1). The sites sampled encompassed the range of streams and gradient of urban impacts across Wellington City. Where practicable, samples were not taken within two weeks of any flood event (flood events are defined as flows greater than three times the median river flow). Samples were collected with the use of a kick-net (0.5 mm mesh size) following Protocol C1 of the national macroinvertebrate sampling protocols (Stark et al., 2001). All samples are processed in accordance with Protocol P2 (Stark et al., 2001).

Macroinvertebrates were collected to calculate the Macroinvertebrate Community Index (MCI) and Quantitative MCI. The MCI is an index of sensitivity to a wide range of environmental variables (Stark and Maxted, 2007) used to measure macroinvertebrate community health. The MCI is used in the GWRC's PNRP to measure ecosystem condition according to river classes defined by Clapcott and Goodwin (2014) (Table 2.2).

Table 2.2: MCI ecological condition classifications from Clapcott and Goodwin (2014) based on river class in the GWRC PNRP

River class		MCI ecological condition class			
		Poor	Fair	Good	Excellent
1	Steep, hard sedimentary	<110	110-120	120-130	≥130
2	Mid-gradient, coastal and hard sedimentary	<80	80-105	105-130	≥130
3	Mid-gradient, soft sedimentary	<80	80-105	105-130	≥130
4	Lowland, large, draining ranges	<90	90-110	110-130	≥130
5	Lowland, large, draining plains and eastern Wairarapa	<80	80-100	100-120	≥120
6	Lowland, small	<80	80-100	100-120	≥120

2.4 Fish

Fish surveys were conducted at 51 sites using backpack electrofishing and/or spotlighting, depending on stream habitat, at representative habitats in each sampling reach. Not all fishing methods were carried out at each site because of dangerous conditions or site access issues. Fishing methods used at each site are listed in Table 2.1. In 2016, inanga spawning assessments were conducted in Kaiwharawhara and Owhiro Streams (see Marshall and Taylor (2018) for more information).

The Index of Biotic Integrity (IBI) was calculated for each site to provide an indication of overall fish community condition and interpretation of scores was based on recommended classes in Joy (2004) (Table 2.3).

Table 2.3: Thresholds for interpretation of IBI scores for the Wellington Region from Joy (2004)

IBI score	Integrity class	Attributes
52–60	Excellent	Comparable to the best situations without human disturbance; all regionally expected species for the stream position are present. Site is above the 97th percentile of Wellington sites.
48–51	Very good	Site is above the 90th percentile of all Wellington sites; species richness is slightly less than best for the region.
38–47	Good	Site is above the 70th percentile of Wellington sites but species richness and habitat or migratory access reduced some signs of stress.
30–37	Fair	Score is just above average but species richness is significantly reduced habitat and or access impaired.
18–29	Poor	Site is less than average for Wellington region IBI scores, less than the 50th percentile, thus species richness and or habitat are severely impacted.
2–17	Very poor	Site is impacted or migratory access almost non-existent.
0	No native fish	Site is grossly impacted or access for fish is non-existent.

2.5 Data analysis

2.5.1 Impervious and vegetated area within each catchment

To get an indication of potential pressure from urban land use, the area and percentage of impervious and vegetated land cover in the catchment upstream of each site was calculated in ArcGIS Pro. Impervious and vegetated areas were defined using methods outlined by Kaspersen et al.(2015) using Sentinel 2b satellite imagery which covers Wellington City on the 15th June 2019. To estimate impervious and vegetated areas the Soil Adjusted Vegetation Index (SAVI) defined by Kaspersen et al.(2015) was used and reclassified so that values less than 0.9 were given the value of 1 (impervious surface) while all others were given 0 (vegetation). The resulting raster dataset was clipped to the Wellington City boundary and the values converted to vector polygons. All analysis was conducted in ArcGIS Pro.

Building outline data and road/rail parcel data were added to the analysis to be joined with the SAVI polygons. Some manual processing of the data also occurred for quality assurance purposes.

2.5.2 Macroinvertebrate and fish community structure

Similarities in macroinvertebrate and fish community structure between sites were analysed using the Bray-Curtis similarity measure, and visualised using non-metric multidimensional scaling ordination (NMDS) in Primer version 7. The Pearson correlation coefficient was calculated to determine the environmental factors correlated with each axis in the NMDS ordination. All analyses were conducted on relative abundance data, which was fourth root transformed.

3. Results

3.1 Habitat condition

Overall, habitat scores ranged from fair (33.5/100 at Ngauranga Stream at Taylor Prestons site N5) to excellent (94/100 at Waipapa Stream site WP01) with a median score of 73.25. Sites with the greatest habitat condition were generally located in stream segments upstream of urban areas (eg. Waimapihi Stream in Aro Valley) or in catchments with no urban cover and regenerating vegetation (eg. Waipapa stream on the Wellington South Coast) (Figure 3.1). A summary of the overall habitat scores and scores for each variable at each site is provided in the supplementary data spreadsheet.




Figure 3.1: Results of habitat assessments at each survey site sampled between 2016 to 2019. The total habitat assessment score is shown for each site.

3.1.1 Macroinvertebrates

MCI scores ranged from 59 (poor ecological condition) at site N1 on Ngauranga Stream at Newlands Road, to 131 (excellent ecological condition) at site N3 on Ngauranga Stream at Mount Kaukau.

The five highest MCI scores were at sites:

- N3 (Ngauranga Stream at Mount Kaukau): 131
- K6 (South Makara Stream Tributary at South Makara Road): 129
- K2 (Karori Stream at Castlemaine Close): 122
- WP02 (Waipapa Stream at South Coast): 122
- WA01 (Waimapihi Stream): 119

The five lowest MCI scores were at sites:

- N1 (Ngauranga Stream at Newlands Road): 58
- N2 (Ngauranga Stream near Alex Moore Park) : 62
- K9 (Karori Stream at Darwin Street): 72
- O4a (Owhiro Stream downstream of Happy Valley Road): 73
- K1 (Karori Stream at Karori Park): 73

The highest MCI scores were generally in areas with the greatest habitat quality (e.g. greater riparian cover) and the lowest levels of urban development in the surrounding catchment (Figure 3.2 and supplementary data spreadsheet). The lowest MCI scores were in catchments with higher levels of urban landuse. As impervious area in the upstream catchment of a site increased there was a decrease in the relative abundance of Ephemeroptera, Plecoptera and Trichoptera (EPT) taxa which are sensitive to poor water and habitat quality. Sites with higher levels of imperviousness in the upstream catchment had higher relative abundances of taxa such as Diptera, Oligochaeta and Mollusca (primarily *Potamopyrgus* and *Physa* species) which are tolerant of poor habitat and water quality (Figure 3.3).

The percentage of impervious area upstream of a site had a possible influence on the type and abundance of macroinvertebrates present at a site, with a high negative correlation between percent impervious area upstream and position on axis 1 in the ordination plot. Furthermore, sites with less impervious area upstream (more vegetation cover upstream) and better habitat condition (e.g. less deposited sediment and greater habitat diversity) were generally associated with greater MCI/QMCI scores (i.e. more taxa sensitive to poor water quality and habitat disturbance), taxa richness, percentage EPT abundance and EPT richness. The amount of riparian shading (assessed using the habitat assessment protocol) also had a high correlation with the composition of the macroinvertebrate community on axis 2 of the ordination (Figure 3.4, Table 3.1).




Figure 3.2: MCI ecological condition class (defined by Clapcott and Goodwin 2014) at each site where macroinvertebrates were collected between 2016 and 2019




Figure 3.3: Relative abundance of macroinvertebrate taxa groups across all sites from 2016-2019. Sites on the x axis are listed in order of increasing impervious area upstream. Impervious area calculations for each site are listed in the supplementary data spreadsheet.




Figure 3.4: NMDS ordination of sites based on fourth root transformed macroinvertebrate relative abundance showing correlations with habitat assessment variables, % imperviousness upstream, % vegetation cover upstream and macroinvertebrate community measures. The closer the line to the circle the higher the correlation. Correlations with each axis are show in Table 3.1.

Table 3.1: Pearson correlations with axes 1 and 2 for variables overlaid in the ordination plot shown in Figure 3.4

Variable	Axis 1	Axis 2
Low deposited sediment	-0.05	-0.67
Invertebrate habitat diversity	0.22	-0.31
Invertebrate habitat abundance	-0.18	-0.66
Riparian shading	0.4	0.56
Total habitat score	0.37	-0.43
Taxa richness	0.81	-0.35
MCI	0.88	-0.17
QMCI	0.82	-0.09
EPT richness	0.84	-0.44
% EPT	0.74	-0.43
% imperviousness upstream	-0.62	0.37
% vegetation upstream	0.62	-0.37

3.1.2 Fish and koura

Fish communities within urban areas generally had fish IBI scores corresponding to fair or poor condition (Figure 3.5). At some sites no native fish were collected (IBI score of 0) and overall IBI scores ranged from 16 (very poor) to 56 (excellent):

The sites with the lowest IBI scores were:

- N5 (Ngauranga Stream at Taylor Prestons): 0
- O2a (Owhiro Stream downstream of TNT Landfill): 0
- O2c (Owhiro Stream at Southern landfill gate on Landfill Rd) : 0
- MH01 (Mahinga Bay Stream) : 0
- PA02 (Island Bay tributary at Mana Karioi) : 0
- TP01 (Te Poti Stream) : 0
- WA02 (Clinical track Stream) : 0
- WT01 (Bells Road Stream) : 0

- K5 (Karori Stream at Makara coast): 16
- Kai4 (Kaiwharawhara Stream below Korimoko confluence): 16
- O2b (Owhiro Stream below Landfill Rd confluence): 16

- K10 (Karori Stream at Futuna Close): 18
- N4 (Ngauranga Stream at Tyres Stream Reserve): 18
- MR01 (Maupuia Stream): 18

The sites with the highest IBI scores were:

- Kaiwharawhara Stream at Spotlight: 56
- WP02 (Waipapa Stream at coast): 50
- K2 (Karori Stream at Castlemaine Close): 42
- PT01 (Puketea Stream at Botanical Gardens): 40
- K6 (South Makara Stream tributary at South Makara Road): 40
- Kai1 (Kaiwharawhara Stream below the dam): 38

The sites with the highest IBI scores all had habitat with good to excellent fish cover diversity and cover abundance (habitat assessment ranging from 8 to 10/10 at sites WP02, K2, PT01, K6 and Kai 1 – see supplementary data). Kaiwharawhara Stream at Hutt Road/Spotlight and Waipapa Stream at the coast had direct access to the sea for migration with no barriers.

In total, fourteen fish species were identified along with koura (Table 3.2). Eel species and banded kokopu were the dominant species across the majority of sites (Figure 3.6). Climbing species such as eels, banded kokopu and koaro were the only species found in open stream sections upstream of pipes (e.g. Kumutoto Stream, Ngaroma Stream, Maupuia Stream and Moturoa Stream). Whereas species with poor climbing ability such as inanga and the bully species were only found in streams with direct access to the sea and no barriers (e.g. lower reaches of the Kaiwharawhara stream) Koura were found in all catchments except Kumutoto, Mahanga and Wadestown (Table 3.2, Figure 3.6). Potential inanga spawning habitat has also been identified and confirmed in the lower reaches of Kaiwharawhara Stream in bank vegetation (Marshall and Taylor, 2018).

The diversity and abundance of fish species was influenced by the diversity of fish habitat available (including riparian shading and hydraulic heterogeneity). The pattern of the fish community composition across sites was correlated with fish habitat cover diversity (Pearson correlation 0.35 axis 1) and riparian shading (Pearson correlation 0.61 axis 1) and hydraulic heterogeneity (Pearson correlation -0.38 axis 2) (Table 3.3, Figure 3.7). As fish habitat cover diversity and riparian shading increased there was a greater range of species present and higher abundance of banded kokopu (Figure 3.6, Figure 3.7). However, the correlations with fish community measures (IBI, taxa richness) and %imperviousness/vegetation cover upstream were not as high as those for the macroinvertebrate community, which indicates there are other key factors affecting fish not accounted for (e.g. fish passage barriers) (Table 3.1, Table 3.3).

The native fish collected included at risk and declining species such as koaro, inanga, redfin bully, longfin eel, giant kokopu (Dunn et al., 2018). The nationally vulnerable shortjaw kokopu was only found in the Waipapa Stream on Wellington's South Coast, which has no urban land use in the surrounding catchment (Table 3.2, Figure 3.6). See the supplementary data spreadsheet for a table of fish species identified at each site. Appendix 1 contains photos from fish sampling from various locations.




Figure 3.5: Fish IBI condition classes at each site where fish were surveyed between 2016 and 2019 (classes defined by Joy (2004) see Table 2.3)

Table 3.2: Summary of fish species (and koura) identified in Wellington urban stream catchments using both electric fishing and spotlighting methods from 2016 to 2019. * Note presence absence results have been added for the Kaiwharawhara Stream at Spotlight from fishing conducted in 2018 and 2019 for the Matariki event.

Catchment	Longfin eel	Shortfin eel	Banded kokopu	Giant kokopu	Shortjaw kokopu	Unidentified kokopu	Koaro	Inanga	Upland bully	Giant Bully	Redfin bully	Black flounder	Triplefin	Koura	Trout	Grey mullet	Unidentified eel	Unidentified galaxiid	Unidentified fish
Kaiwharawhara*	22	3	107	1		1				1	6	1	1	13			22		
Kumutoto			26				14												
Owhiro	54	15	15					7			14			3	2	1	1		
Karori	54	2	15	1			29		33					14	1		24		
Ngauranga	35	1	7				5	15						2			10		
Motoroa			5				6							2					
Lyll Bay	4	1	40											20					
Mahanga																	1		
Miramar		1	50											4					
Paikawakawa			12											22					
Pipitea	4		9				4							6			4		3
Te Poti														1			1		
Waimapihi			17				5							103				4	5
Waipapa	23	4	5	1	1		22							1				1	
Waitangi			6											28					
Wadestown	1		10																




Figure 3.6: Relative abundance of fish species (and koura) identified in Wellington urban stream catchments using both electric fishing and spotlighting methods from 2016 to 2019. * Note results are not shown for the Kaiwharawhara Stream at Spotlight because this is presence absence data for fishing conducted in 2018 and 2019 for the Matariki event. Sites on the x axis are listed in order of increasing fish cover diversity measured by the habitat condition assessment. Fish cover diversity assessments were not undertaken at all sites.




Figure 3.7: NMDS ordination sites based on fourth root transformed fish relative abundance showing correlations with habitat assessment variables, % imperviousness upstream, % vegetation cover upstream and fish community measures. The closer the line to the circle the higher the correlation.

* Note results are not shown for the Kaiwharawhara Stream at Spotlight because this is presence absence data for fishing conducted in 2018 and 2019 for the Matariki event.

Table 3.3: Pearson correlations with axes 1 and 2 for variables overlaid in the ordination plot shown in Figure 3.6

Variable	Axis 1	Axis 2
Low deposited sediment	-0.27	-0.29
Fish cover abundance	-0.07	0.18
Fish cover diversity	0.35	0.24
Riparian shading	0.61	0.22
Total habitat score	0.21	-0.07
Fish taxa richness	-0.22	-0.04
IBI	0.25	-0.01
Increased hydraulic heterogeneity	-0.0003	-0.38
% imperviousness upstream	-0.07	0.04
% vegetation upstream	0.07	-0.04

4. Discussion and conclusions

4.1 Urban stream ecosystem health and implications for urban stream management in Wellington City

The ecosystem health of urban streams within Wellington City (as represented by macroinvertebrate communities) is variable. Factors such as habitat diversity, riparian shading and impervious area in the upstream catchment influence the ecological health of Wellington City urban streams. The surveys conducted to date as a part of the Wellington City urban streams monitoring programme have provided information which can be used to set a baseline for the current state of ecosystem health values related to habitat, macroinvertebrates and fish.

Macroinvertebrate community health was best at Ngauranga Stream at Mt Kaukau, South Makara Stream tributary at South Makara Road and Karori Stream at Castlemaine Close, and worst at Ngauranga Stream at Newlands Road, Ngauranga Stream near Alex Moore Park and Karori Stream at Darwin Street. Macroinvertebrate communities were generally in better condition (as indicated by the greater presence of sensitive EPT taxa) in streams with less urban landuse and impervious area in the upstream catchment (Figure 3.3). Streams with urban areas upstream were likely to have more deposited fine sediment and lower quality habitat for macroinvertebrates. This result is in line with previous research on urban streams within the Wellington City area which showed there was declining ecosystem health indicated by macroinvertebrate communities when impervious area upstream increased (Warr, 2009). Increased impervious area in urban catchments can result in shorter and more severe high flow events, which can lead to stream bank erosion/sedimentation and effect macroinvertebrate communities in urban streams (Walsh et al., 2005).

Wellington City urban streams also support at risk and declining fish species such as koaro, inanga, redfin bully, longfin eel and giant kokopu (Dunn et al., 2018). The nationally vulnerable shortjaw kokopu (Dunn et al., 2018) was only found in the Waipapa Stream catchment, which has no urban landuse, emphasising that changes to stream habitat from urbanisation may have influenced the distribution of this species within Wellington City.

The condition of native fish communities within the streams assessed ranged from poor to excellent (Figure 3.5). The lowest IBI scores were at sites with no native fish collected in the Ngauranga, Owhiro, Mahinga Bay, Paikawakawa, Te Poti, Waimaphi and Waitangi Stream catchments. The highest fish community condition assessed using the IBI was at Kaiwharawhara Stream at Spotlight (excellent), Waipapa Stream at Coast (very good) and Karori Stream at Castlemaine Close (good). The diversity and abundance of fish species was influenced by factors such as habitat diversity, direct connection to the sea (Figure 3.6) and spawning habitat (e.g. inanga spawning in lower Kaiwharawhara Stream bank vegetation – Marshall and Taylor (2018)). However, not all of the habitat and environmental data collected explained the variation in fish community composition (Table 3.3). This means there are other factors which are influencing fish community composition. For example, barriers to fish movement are one of the major unmeasured pressures

influencing native fish communities in Wellington urban streams. For example, the only fish species found upstream of piped sections of stream were climbing species such as banded kokopu, koaro and eel species. The presence of banded kokopu and eels in piped stream sections was confirmed in the piped stream survey (Figure 3.6, Appendix 2).

From the streams sampled as part of this programme there were generally different pictures of macroinvertebrate and fish community condition. For example, most sites with good macroinvertebrate community health were higher in the catchment and had poor fish community health. The stream sites with best fish community health tended to be in the lower reaches of streams where connectivity to the sea was best. Macroinvertebrate communities are strongly driven by urban activities in the catchment upstream that affect habitat and water quality (indicated by the percent cover for impervious area upstream). In contrast, fish communities appear to be more strongly affected by activities in the downstream catchment that affect fish migration (eg. fish barriers). Both macroinvertebrate and fish communities were related to riparian shading, which could be having an influence through reducing stream temperature or organic matter inputs. These differing relationships with environmental variables need to be taken into account when considering management actions to improve ecosystem health in Wellington City streams.

The initial piped stream survey has shown that some sections of piped streams in Wellington City have habitat which supports fish and macroinvertebrate life (see Appendix 2 reports). For example:

- Some sites have natural cobble and gravel substrates (overlying a concrete base, likely transported from upstream remnant open stream sections).
- Fish appear to be widespread in the piped stream network and were recorded at five of the six survey locations. However, diversity is low and is dominated by eels. In contrast, open stream sections tend to be dominated by banded kokopu. Inanga were also found in piped sections with direct access to the ocean. However, without spawning habitat in the tidal zone of the pipes it is highly unlikely they will ever be able to successfully spawn.
- Based on initial macroinvertebrate samples (there are 72 surber samples still to be analysed by late 2019), of the five most abundant macroinvertebrate taxa in both open and piped streams, three were the same – *Potamopyrgus* and *Physa* snails and Oligochaete worms. These taxa are more tolerant of poor habitat and water quality conditions.
- In open stream sites there was greater taxa richness and a higher percentage of sensitive EPT taxa than in piped stream sites. This is likely to be indicative of better habitat conditions at open stream sites. The complete enclosing of open channels significantly reduces the quality of habitat for aquatic life and results in only a subset of taxa within the catchment being able to persist there.

Further recommendations for the future management of piped streams and future monitoring options will be provided in a final report in early 2020.

4.2 Next steps for Wellington urban stream monitoring

The next stage of the Wellington City urban stream monitoring programme will focus on identifying any remaining streams which have not been sampled. These include streams within Wellington City peri-urban areas which may be subject to future development (Makara and Ohariu) and streams in the part of WCC located in the Porirua Harbour catchment. These streams will be the focus of monitoring in 2019/20.

In addition, a long term monitoring programme for Wellington City streams (stage two) will be designed in collaboration with WCC in 2019/20 for implementation in 2020/21. Aspects which should be considered as part of the monitoring programme design include:

- Agreement between GWRC and WCC urban ecology and policy staff on the objectives of the monitoring programme. Currently the main objective of the programme is to provide information to report on state and trends in ecosystem health within Wellington City urban streams. If the monitoring programme will inform policies and strategies this will mean establishing in the planning stage which questions the programme will and will not be able to answer, and questions which will need to be addressed by targeted investigations or other means.
- A coverage of sites which takes into account the range of locations and ecological/biodiversity values from the sites sampled to date. These sites should include sites with minimal urban impact, sites within the town belt in urban Wellington and sites within the urban areas with piped stream areas upstream/downstream. Having a range of sites with a varying degree of urban impacts such as impervious surfaces and stream piping will enable to effect of urbanisation on fish and invertebrates to be tracked through time.
- Opportunities for additional sites monitored by community citizen science groups to compliment the GWRC collected dataset.
- Additional, less frequent, inanga spawning habitat assessments.
- Ongoing piped stream sampling, subject to final recommendations in 2020.
- Type and frequency of reporting required.

Other knowledge gaps to address as part of the design of the monitoring programme or targeted studies include:

- Assessment of barriers to fish movement throughout the city.
- Assessments of fish population structure. It is currently not known how well fish populations in Wellington urban streams are recruiting.

- Assessments of food web and ecological processes as drivers of ecosystem health.
- Understanding macroinvertebrate community responses to stressors using stressor specific metrics (Clapcott et al., 2017).

Acknowledgements

Stage 1 of this project was jointly funded by Wellington City Council. Thanks to Daniela Biaggio, Michele Frank and Myfanwy Emeny from WCC for their assistance. Thanks to Summer Greenfield for peer reviewing this report and providing many constructive comments. Field surveys were conducted by GWRC staff (Shyam Morar, Emily Martin, Bryn Hickson Rowden, Ashely Alberto Ashley Mitchell, Megan Oliver, Mark Heath and Lucy Baker) and volunteers (Dylan Kulik). The piped stream surveys were conducted by Alex James from EOS Ecology with assistance from Silver Linings Contracting.

References

- Clapcott J, Wagenhoff A, Neale M, Storey R, Smith, B, Death R, Harding J, Matthaeci C, Quinn J, Collier K, Atalah, J, Goodwin E, Rabel H, Mackman J and Young R. 2017. *Macroinvertebrate metrics for the National Policy Statement for Freshwater Management*. Cawthron Report No. 3073 prepared for Ministry for the Environment.
- Clapcott J.E. 2015. *National rapid habitat assessment protocol development for streams and rivers*. Cawthron Institute Report No. 2649 prepared for Northland Regional Council.
- Clapcott JE and Goodwin E. 2014. *Technical report of Macroinvertebrate Community Index predictions for the Wellington Region* Cawthron Institute Report No. 2503 prepared for Greater Wellington Regional Council.
- Dunn N, Allibone R, Closs G, Crow S, David BO, Goodman, JM., Griffiths M, Jack D, Ling N, Waters J and Rolfe J. 2018. *Conservation status of New Zealand freshwater fishes, 2017*. Department of Conservation.
- James A. 2015. *Lambton Harbour Catchment ICMP Stage 2: Ecological Assessment*. EOS Ecology Report No. WEL02-15026-01 prepared for Wellington Water Limited
- Joy M. 2004. *A fish index of biotic integrity for the Wellington region*. Report prepared for Greater Wellington Regional Council.
- Kaspersen P, Fensholt R and Drews M. 2015. *Using Landsat Vegetation Indices to Estimate Impervious Surface Fractions for European Cities*. Remote Sens. 7, 8224–8249.
- Marshall W and Taylor M. 2018. *Inanga spawning habitat surveys in the Wellington Region, 2017*. AEL Report No. 146 prepared for Greater Wellington Regional Council.
- Mitchell A and Heath M. 2019. *Rivers Water Quality and Ecology monitoring programme, Annual data report, 2017/18*. Greater Wellington Regional Council Report No. GW/ESCI-T-18/142, Wellington.
- Stark JD, Boothroyd I, Harding J, Maxted JR and Scarsbrook M. 2001. *Protocols for Sampling Macroinvertebrates in Wadeable Streams*. Ministry for the Environment.
- Stark JD and Maxted JR. 2007. *A user guide for the Macroinvertebrate Community Index*. Cawthron Institute Report No. 1166 prepared for Ministry for the Environment.
- Walsh CJ, Roy AH, Feminella JW, Cottingham,PD, Groffman,PM and Morgan, RP. 2005. *The urban stream syndrome: current knowledge and the search for a cure*. J. North Am. Benthol. Soc. 24, 706–723.
- Warr S. 2009. *Development of an ecological classification for urban streams in the Wellington region*. Greater Wellington Regional Council Memo No. ENV/05/03/14.

Appendix 1: Photos from fish sampling



Figure A1.1: Banded kokopu and a koaro (bottom left) from Moturoa Stream in Central Park. Photo S. Morar



Figure A1.2: Fishing in resident's yards on Ranelagh Street in Karori – longfin eels, shortfin eels and koura. Photo S. Morar



Figure A1.3: Karori Stream on the south coast - Lots of small upland bullies, koura and longfin eels. Photo S. Morar



Figure A1.4: Koaro from a tributary of the Karori Stream. Photo S. Morar



Figure A1.5: Ngauranga Stream 400m above mouth to Wellington Harbour - longfin eels and inanga. Photo S. Morar



Figure A1.6: Ngauranga Stream at end of Tyers Road off Centennial Highway - lots of koaro and koura. Photo S. Morar



Figure A1.7: A large giant kokopu from Kaiwharawhara Stream near the Old Hutt Road Bridge at Spotlight. Photo S. Morar



Figure A1.8: Koaro in Kumutoto Stream at Victoria University. Photo S. Morar



Figure A1.9: Whitebait sized banded kokopu from Ngaroma Stream in Lyall Bay. Photo S. Morar



Figure A1.10: Koura from Farnham Street Stream. Photo S. Morar

Appendix 2: Freshwater Ecology of Piped Streams in Wellington: Pilot Study Interim and Final Reports



Freshwater Ecology of Piped Streams in Wellington: Pilot Study Interim Report

EOS Ecology Report No. GRE01-17087-01 | July 2019

Prepared for Greater Wellington Regional Council

Prepared by EOS Ecology – Alex James

Reviewed by Shelley McMurtrie (EOS Ecology)



www.eosecology.co.nz
info@eosecology.co.nz

PO Box 4262, Christchurch 8140
P 03 389 0538

PO Box 8054, Palmerston North 4446
P 06 358 9566

SCIENCE + ENGAGEMENT



EOS Ecology or any employee or sub-consultant of EOS Ecology accepts no liability with respect to this publication's use other than by the Client. This publication may not be reproduced or copied in any form without the permission of the Client.

© All photographs within this publication are copyright of EOS Ecology or the credited photographer; they may not be used without written permission.

1	INTRODUCTION.....	1
2	METHODS.....	2
2.1	Health and Safety.....	2
2.2	Site Selection.....	2
2.3	Detailed Ecological Survey.....	5
3	RESULTS.....	7
3.1	Habitat.....	7
3.2	Fish.....	10
3.3	Benthic Macroinvertebrates – Manhole Lifting Kick Nets.....	12
4	CONCLUSIONS.....	15
4.1	Habitat.....	15
4.2	Fish.....	15
4.3	Benthic Macroinvertebrates – Manhole Lifting Kick Nets.....	15
5	MEDIA.....	16
6	ACKNOWLEDGEMENTS.....	16
7	REFERENCES.....	16

1 INTRODUCTION

In most cities urban streams have been extensively piped to provide land for development and alleviate flood and disease risk. In many catchments open stream habitat exists only as short remnant channels. To date ecological information on these highly modified catchments has primarily been derived from these remnant open sections for reasons of accessibility and safety, even though these sections make up a small percentage of total stream length. Further, the piped sections that join isolated open reaches are often managed as part of the stormwater network and have typically been ignored from an ecological perspective.

Wellington, with its steep topography and coastal location is a good example of a city where numerous small coastal stream catchments have been extensively piped, such that in several suburbs open stream channels are now only found as fragmented remnants. With the exception of some anecdotal reports of eels down stormwater grates, piped streams in Wellington have previously only been considered as migration pathways for some freshwater fishes that are known to be present in remnant open sections (e.g. banded kokopu, koaro, eels). The fish and macroinvertebrates living within the piped sections are unknown and there has never been any attempt to characterise freshwater habitat condition in these highly modified stream environments.

EOS Ecology undertook a series of urban stream catchments investigations for Wellington Water Limited (WWL) to support Integrated Catchment Management Plan (ICMP) development from 2015 to 2017, where the project brief was only to examine open sections of streams. It became apparent the ecological values of extensively piped catchments could not be fully determined without examining the piped sections (which were often the higher proportion of total stream length). Based on the catchment knowledge gained during the ICMP investigations EOS Ecology then developed a plan to undertake a pilot study to survey piped stream ecology in Wellington. The lack of information on piped stream ecology was suggested to Greater Wellington Regional Council (GWRC) as a major knowledge gap and they agreed to fund a pilot study of six sites focussing on fish, macroinvertebrates, and habitat quality.

EOS Ecology was contracted by GWRC to design, implement, and report on this pilot study. This interim report includes methodology, fish data, and manhole macroinvertebrate data; with a later final report to come once the detailed survey macroinvertebrate data is available. The final report will compare macroinvertebrate data among sites, catchments, method (manhole kick net vs in pipe Surber sampling), and between open and piped streams. It will also recommend future monitoring and identify knowledge gaps.

2 METHODS

2.1 Health and Safety

Underground pipes are classified as confined spaces meaning only appropriately trained persons are able to enter and special procedures are required (e.g., operating a permitting system, use of gas detectors, use of a winch and rescue-tether system). As WWL manages the piped stream network as part of the stormwater system, a WWL-approved contractor was required to assist with the project. WWL put us in touch with Silver Linings Contracting, who were then contracted to assist with all piped stream fieldwork. They were in charge of site health and safety and supplied all equipment required for confined space entry. From EOS Ecology, the author (Alex James) was confined spaces entry certified.

2.2 Site Selection

2.2.1 Desktop Exercise

WWL provided GIS layers of the stormwater network, manhole locations, and remnant open channels. These were overlain with Google Earth imagery and used to select manholes within Wellington's inner suburbs where entry for an ecological survey appeared logistically realistic. Criteria for potential sites including:

- » Pipe diameters of no less than 1200 mm.
- » Manholes to be located on footpaths and berms to avoid the requirement for traffic management or entering of private property (Figure 1).
- » The catchment would preferably have permanently flowing open stream remnants from which fish were known.

Based on these criteria, 36 candidate manholes were identified with the expectation some would be inaccessible in the field.



Miramar Stream manhole on grassed berm



Pae Kawakawa Stream manhole on footpath

Figure 1 Examples of berm and footpath locations of candidate manholes.

2.2.2 Manhole Lifting Exercise

Over 1–2 May 2018 a manhole lifting site visit was undertaken with Silver Linings Contracting to aid selection of the final survey sites. Overall 20 manholes were opened with the remaining 16 either being inaccessible (e.g., buried, car parked over, building on top) or being in close proximity to an opened manhole. We also took the opportunity to trial macroinvertebrate sampling from the surface utilising a modified sampling net.

Once each manhole was opened a set procedure was undertaken:


- » Lowering of a gas detector to check for unsafe concentrations of oxygen, hydrogen sulphide, carbon monoxide, and flammable gases (Figure 2).
- » Measurement of surface to pipe bottom and pipe diameter using a Leica Disto laser measurement device.
- » Visual estimation of water depth (dry/shallow/medium/deep), substrate type (brick, concrete, silt, cobbles, other), pipe shape/profile (circular/rectangular/arch/other), pipe material (concrete/brick/other).
- » Collection of macroinvertebrate samples using a brush and standard kick net with extendable handles (2.7–5 m). Such samples were collected from 16 sites (Figure 2). Samples were preserved with isopropyl alcohol (IPA) for later processing.
- » The taking of photos of the manhole shaft and stream below.
- » The taking of video footage from selected manholes using a GoPro video camera and torch attached to the extendable handle kick net (Figure 2).




Using the extendable-handle kick net to collect a macroinvertebrate sample



Using the extendable-handle kick net in conjunction with a brush to collect a macroinvertebrate sample



Measuring the atmosphere using a gas detector



The GoPro video camera setup

Figure 2 Examples of manhole lifting exercise methodology.

2.2.3 Final Site Selection

Six sites were selected for detailed ecological survey; three in Pae Kawakawa Stream catchment (Island Bay); two in Miramar Stream (Miramar); and one in Waipapa Stream (Hataitai) (Figure 3). The Pae Kawakawa Stream and Miramar Stream sites were all vertical entries down manholes, while Waipapa Stream involved entry via the ocean outlet at low tide. It was originally planned to have a site in the Waitangi Stream catchment, however the only suitable site was deemed too dangerous to enter without using breathing apparatus, for which extra specialist training would have been required. A third Pae Kawakawa Stream site was substituted.



Figure 3 Locations of the six detailed survey sites.


2.3 Detailed Ecological Survey

The six sampling sites were visited on two occasions for the detailed survey (stage 1: 6–7 March 2019 and stage 2: 20–21 March 2019). Stage 1 involved installation of trail/game cameras, sticky traps, and collection of macroinvertebrate samples, while stage 2 consisted of recovering of cameras and sticky traps and undertaking a fish survey. In practice fish were often sighted during stage 1, so an informal fish survey (consisting of noting if fish were seen) was also undertaken at that time. Additionally, an equipment mishap required macroinvertebrates at one site to be collected on the stage 2 visit. Details of the survey methodology are detailed below:

- » **Trail/game cameras:** At each site a single trail camera was installed near the manhole with the lens facing towards the wetted channel in an attempt to obtain images of fish within the piped streams (Figure 4). Cameras were mounted on existing pipes or reinforcing steel rods where available, otherwise a steel extendable curtain rod was affixed across the width of the piped stream. Curtain rods were attached using cable ties to any available protrusions (e.g., ladder rungs, reinforcing steels) and/or appropriate concrete adhesives applied via caulking gun. Cameras were set to take time-lapse images every 10 minutes for the duration of deployment and also be triggered to take a photo by movement. The typical movement sensor of such cameras relies on a difference in temperature between the environment and the animal, hence, is unlikely to be triggered by fish. Where the camera model allowed (we used three different models) 10-second videos were also recorded every 10 minutes.
- » **Sticky traps:** At each site a single 24.5 cm wide by 40 cm high plastic yellow sticky trap was installed as near the roof of the piped stream as possible (Figure 4). These were sticky on both the upstream and downstream sides and installed to existing pipes or reinforcing steels at some sites or to the curtain rod described above at others.
- » **Macroinvertebrates:** Twelve Surber samples (0.09 x 0.09 m²) were collected from each site over a 40 to 100 m length of stream depending on availability of habitat suitable for Surber sampling and access (Figure 4). Samples were preserved with isopropyl alcohol (IPA) for later processing.
- » **Fish survey:** At each site a 200 m (100 m upstream and downstream of the manhole access point) reach was carefully searched for fish using a spotlight and hand netting technique (Figure 4). The exception was Waipapa Stream where we entered via the ocean outlet. At this site we searched for fish from the entrance all the way upstream to where the macroinvertebrate survey was undertaken. In practice in Waipapa Stream fish were only observed in the lower 90 m of pipe as above this point water became too shallow and cover lacking for the fish present. Any obvious fish cover elements (e.g., pieces of wood, brick, and assorted rubbish) were slowly lifted to determine if fish were underneath. Fish were identified to species, with the exception of those eels that were unable to be captured. Lengths were visually estimated.
- » **Habitat characterisation:** At each Surber sample location (12 at each site) water depth, wetted width, substrate composition, and organic matter cover (leaves, CPOM, biofilms) was recorded. We recorded the “head” where water depth was recorded to enable an estimate of water velocity using the ruler methodology (cf. Harding et al., 2009 – Appendix 3). Biofilm colour and thickness was also recorded. Site photos were taken including images of each Surber location prior to sampling (Figure 4). We also made site wide measurements of meso-habitat length (rapid, run, riffle, pool, fall) and made visual estimates of substrate composition and organic matter cover (leaves, roots, CPOM, biofilms). Any potential fish barriers were noted.
- » **Water quality:** A YSI multiprobe supplied by GWRC was used to collect spot water quality records of temperature, dissolved oxygen, and conductivity.




Camera and sticky trap attached to reinforcing steel



Surber sampling



An eel captured via spotlighting and hand netting



Example of substrate sampled for macroinvertebrates

Figure 4 Images of detailed survey methodologies.

3 RESULTS

3.1 Habitat

Water depths were generally very shallow with only one site (the tidally influenced Waipapa Stream) having an average greater than 0.1 m (Table 1). Two sites had relatively swift water velocities (The Parade – Dover St and Waipapa), which were a result of pipe gradient. The other sites had very low water velocity with the 302 The Parade and Miramar – Shops sites having no detectable water velocity with the velocity head rod methodology used (Table 1). Two sites had virtually no mobile substrate with the streambed being entirely bare concrete (The Parade – Dover St) or bricks (Waipapa) (Table 1, Figure 5). These were also the two sites with the relatively high water velocities; hence it appears there are minimal depositional areas in these high gradient sections. The 302 The Parade site was also predominantly concrete but did have some areas of deposited substrate. The three other sites had a significant cover of a range of stony and sandy substrates (Table 1, Figure 5). Biofilms were prominent at all sites with the exception of 348 The Parade where there were none detected (Table 1). Other organic matter such as leaves and wood were relatively uncommon. Mesohabitat was predominantly runs, with the only riffle-type habitat present at 348 The Parade. Deeper pool habitat was only present at 302 The Parade and 348 The Parade (Table 1).

Spot water temperatures were in the 16.4–18.5 °C range across the sites, while specific conductivity was very high (28,440 $\mu\text{S}/\text{cm}$) at the Miramar – Shops site due to the influence of seawater (Table 1). The other five sites had relatively high specific conductivities (323–481 $\mu\text{S}/\text{cm}$) for freshwater systems. All sites had well oxygenated water with only the tidally influenced Miramar – Shops site being less than 80% saturation at the time of measurement (Table 1).

Table 1 Habitat characteristics measured at each Surber sampling location (n=12 per site), mesohabitat percentages measured over the length of the macroinvertebrate sampling reaches, and water quality spot measures recorded at the entrance manholes (or start of macroinvertebrate sampling reach in Waipapa Stream).

Parameter	Pae Kawakawa Stream			Waipapa Stream	Miramar Stream	
	The Parade – Dover St	302 The Parade	348 The Parade	Waipapa*	Miramar Park	Miramar – Shops**
Mean wetted width (m)	1.17	0.63	1.14	0.52	1.25	1.38
Mean water depth (m)	0.07	0.05	0.05	0.07	0.04	0.16
Mean water velocity (m/s)	0.51	0	0.05	0.61	0.02	0
Mean substrate composition (%)	Concrete: 100	Concrete: 77 Cobble: 2 Pebble: 3 Gravel: 5 Sand: 13	Concrete: 12 Cobble: 3 Pebble: 4 Gravel: 39 Sand: 41	Brick: 100	Cobble: 5 Gravel: 13 Pebble: 80 Sand: 2	Concrete: 25 Cobble: 5 Pebble: 37 Gravel: 31 Sand: 2
Mean organic matter cover (%)	Biofilms: 100 Leaves: 0.1	Biofilms: 76 Wood: 0.2 Leaves: 0.7	Wood: 2 Leaves: 1	Biofilms: 90	Biofilms: 92.5 Wood: 0.2 Leaves: 0.6	Biofilms: 60
Mesohabitat lengths (%)	Run: 100	Run: 97 Pool: 3	Run: 51 Riffle: 44 Pool: 5	Run: 100	Run: 100	Mesohabitat varies with tide
Spot temperature	16.4 °C	18.5 °C	18.2 °C	17.4 °C	17.7 °C	17.8 °C
Spot specific conductivity	481 µS/cm	323 µS/cm	580 µS/cm	371 µS/cm	440 µS/cm	28,440 µS/cm
Spot dissolved oxygen	83.6% 8.27 mg/L	93.4% 8.86 mg/L	83.6% 7.95 mg/L	95.2% 9.22 mg/L	84.6% 8.16 mg/L	74.8% 6.49 mg/L
Spot water quality measurement time	10:30 am	4:15 pm	3:00 pm	12:20 pm	9:00 am	2:15 pm

*The Waipapa site data presented was collected in the reach where macroinvertebrates were sampled, all fish were observed in the lower, tidally influenced 90 m of pipe, which had a mostly bare concrete base.

**The Miramar – Shops site was tidally influenced hence wetted width and depth will vary over tidal cycles.



The Parade – Dover St



302 The Parade



348 The Parade



Waipapa Stream



Miramar Park



Miramar - Shops

Figure 5 Images of each site within the macroinvertebrate sampling reach.

3.2 Fish

Fish were present at five of the six sites that underwent detailed ecological survey, with only The Parade – Dover St site not having any fish found over the 200 m survey reach (Figure 6). A total of 54 fish were found at the five piped sites. Fish abundance ranged from 6–14 individuals with the 348 The Parade site having the highest number of fish (15 fish) and the 302 The Parade site having the least (six fish) (Figure 6). Species diversity was low with four species being found across the sites, which were (in order of abundance) Longfin eel, shortfin eel banded kokopu, and inanga (Figure 6). Longfin and shortfin eels were found at all sites where fish were found, whilst two banded kokopu (a recently dead adult and a post-whitebait juvenile) and one inanga were only found at one site each (Figure 6, Figure 7). The two kokopu were found at the Miramar Park site. One was a recently dead adult banded kokopu (150 mm long), which based on the markings on the body had been killed by a large eel (Figure 6). One banded kokopu post-whitebait juvenile (50 mm long) was also found at this site. The single inanga was a young adult (70 mm) found at the tidally influenced Miramar – Shops site (Figure 6, Figure 7).




Figure 6 Fish community metrics and relative abundances from the six piped stream sites surveyed on 20–21 March 2019. 'UnIDed eel' = unidentified eels, and refers to those eels that were not able to be caught to allow for a definitive identification.



Longfin eel



Inanga young adult



Post-whitebait juvenile banded kokopu



Banded kokopu adult (dead)

Figure 7 Fish species found during surveys of the six piped stream sites on 20–21 March 2019.

For the Pae Kawakawa and Miramar Stream catchments there was recent GWRC fish survey data available from open stream sites allowing for comparison of fish assemblages between remnant open stream sections and piped stream sections (Figure 8). In Pae Kawakawa Stream no fish (or waikoura) were found at one open and one piped site. At sites where fish were found there was a distinct difference in composition between open and piped stream sections. Shortfin and longfin eel were found exclusively at the piped sites while the open site with fish had banded kokopu (Figure 8). In Miramar Stream eels (shortfin and longfin) also dominated in the piped sites and longfin eels were only found at piped sites, while the single open stream site surveyed had mostly banded kokopu (Figure 8). Waikoura (freshwater crayfish) were only found in open stream sites in both catchments. Overall, the addition of the piped stream surveys has increased the known diversity of fish in both catchments by two species (longfin and shortfin eel for Pae Kawakawa Stream, and inanga and longfin eel for Miramar Stream).




Figure 8 Relative abundance of fish species (including waikoura) at piped and open survey sites in the Pae Kawakawa Stream and Miramar Stream catchments. Open stream sites were surveyed by GWRC.

3.3 Benthic Macroinvertebrates – Manhole Lifting Kick Nets

A total of 21 invertebrate taxa were recorded from the 16 sampled manholes. The most diverse groups were the two-winged flies (Diptera: 6 taxa), molluscs (Mollusca: 5 taxa), and crustaceans (Crustacea: 3 taxa). Groups represented by one taxon included caddisflies (Trichoptera), springtails (Hexapoda: Collembola), mites (Arachnida: Acari), and four groups collectively called worms (Nematoda, Nemertea, Oligochaeta, Platyhelminthes).

The macroinvertebrate data collected during the manhole lifting exercise were compared to a GWRC dataset from the recent sampling of urban stream sites in Wellington city. This GWRC open stream sampling was undertaken in April 2018 and for comparison with our piped stream data we included sites within Karori Stream (three sites), Kaiwharawhara Stream (one site), Moturoa Stream (Central Park; two sites), Ngauranga Stream (two sites), and Papawai Stream (Prince of Wales Park; one site).

Potamopyrgus snails and Oligochaeta worms were the most abundant taxa overall in both open and piped urban stream sites, albeit in different order (Figure 9). Of the five most abundant taxa open stream and piped stream sites had three in common (*Potamopyrgus*, Oligochaeta worms, and *Physa* snails) (Figure 9). The only insect taxon among the most common taxa in both open stream and piped stream sites were Chironomidae midge larvae – Orthocladiinae in open streams and Chironominae/*Polypedilum* in piped streams (Figure 9).

Open streams had greater taxa richness and percentage of pollution-intolerant EPT taxa (Figure 10). Open streams also had higher MCI scores with an overall mean just inside the “fair” interpretative category of Stark & Maxted (2007). The piped stream overall mean MCI was well within the “poor” category (Figure 10). Relative abundances of higher taxonomic groupings were similar with molluscs, oligochaetes, and Diptera being the most prominent groups in open

and piped streams. Oligochaete worms were however particularly abundant at the piped stream sites, accounting for around 50% of all animals captured (Figure 10).











Open streams	Piped streams
 <p data-bbox="347 656 655 685"><i>Potamopyrgus</i> snails (29%; MCI=4)</p>	 <p data-bbox="911 656 1219 685">Oligochaeta worms (50%; MCI=1)</p>
 <p data-bbox="354 931 649 960">Oligochaeta worms (22%; MCI=1)</p>	 <p data-bbox="904 931 1216 960"><i>Potamopyrgus</i> snails (20%; MCI=4)</p>
 <p data-bbox="320 1207 689 1236">Orthoclaadiinae midge larvae (12%; MCI=2)</p>	 <p data-bbox="948 1207 1179 1236"><i>Physa</i> snails (9%; MCI=3)</p>
 <p data-bbox="384 1482 616 1512"><i>Physa</i> snails (11%; MCI=3)</p>	 <p data-bbox="890 1482 1230 1512"><i>Polypedilum</i> midge larvae (8%; MCI=3)</p>
 <p data-bbox="341 1758 668 1787"><i>Paracalliope</i> amphipods (6%; MCI=5)</p>	 <p data-bbox="954 1758 1169 1787">Collembola (5%; MCI=6)</p>

Figure 9 The five most abundant taxa in Wellington open urban streams (9 sites combined) and pipes streams (16 sites combined). The relative abundance percentages and MCI-hb scores for each taxon are shown in parentheses. All images © EOS Ecology except *Polypedilum*, which is by Landcare Research.




Figure 10 Macroinvertebrate community metrics and relative abundance comparisons between Wellington open urban streams (9 sites combined) and piped streams (16 sites combined). Piped stream sites only includes data from the kick net samples collected during the initial manhole lifting exercise. %EPT is based on abundance data (not taxa richness). Error bars are one standard error. For MCI-hb the generic nationwide water quality interpretative categories of Stark & Maxted (2007) are shown. Wellington-specific MCI classes were not used as GWRC’s river classification does include a class for piped streams.

4 CONCLUSIONS

4.1 Habitat

- » Some sites have natural cobble and gravels substrates (usually overlaying a concrete base), which have presumably been sourced from upstream remnant open stream sections and transported downstream by high flows. Other sites had minimal or no loose rocky substrates. It is likely that piped streams have depositional and erosional zones based on gradient and water velocity as in natural stream channels.
- » Water depths are generally very shallow (<10 cm) and deeper pool habitat is generally very limited or absent.
- » Brown-orange biofilms were prominent at five of the six sites and presumably are a food source for some macroinvertebrates.
- » Spot measures of dissolved oxygen indicated relatively well oxygenated water.

4.2 Fish

- » Fish appear to be widespread in the piped stream network, being present at five of the six detailed survey sites.
- » Piped stream fish diversity is low and dominated by eels. Both longfin and shortfin eels are present.
- » Inanga have the ability to live within piped streams. In catchments that are piped to the ocean it is highly unlikely they could ever successfully spawn due to a lack of spawning habitat.
- » The eel population and lack of cover likely contributes to the general rarity of galaxiid species within piped streams.
- » There appears to be clear habitat partitioning of the fish fauna (and waikoura) in catchments that have extensive piped sections and remnant open headwater sections. Eels dominate the pipes while banded kokopu (and also waikoura) dominate the open sections.
- » The resident eels likely feed intensely on upstream migrating whitebait and elvers at certain times of the year.

4.3 Benthic Macroinvertebrates – Manhole Lifting Kick Nets

- » It is possible to obtain macroinvertebrate samples from manholes without entering the piped system, thus avoiding a confined space entry. However, in practice collecting samples with a standard kick net on an extendable handle was awkward, especially down deeper shafts, and only allowed a very small area of streambed to be sampled. Comparison of manhole samples with detailed survey Surber samples will determine if manhole samples are sufficient to characterise the macroinvertebrate community of piped streams. This will be undertaken later in this Pilot Study.
- » Based on manhole samples the macroinvertebrate assemblage of piped streams was quite similar to that off open urban streams with *Potamopyrgus* snails and oligochaeta worms being the two most abundant taxa at both site types. Of the five most abundant taxa, three were common between open and piped stream sites (*Potamopyrgus* snails, oligochaeta worms, and *Physa* snails).
- » Open stream sites had much greater taxa richness and higher %EPT taxa than piped stream sites.
- » Catchment urbanisation results in a series of environmental filters, which cause the decline and disappearance of a number of freshwater invertebrate taxa. Complete enclosing of most of the open channels in a catchment adds a further severe filter and creates conditions in which only a small subset of available taxa are able to persist.

5 MEDIA

» Ecological work in piped streams has the ability to capture the imagination of journalists as a press release by GWRC prior to the fieldwork resulted in stories by TVNZ's One News and Radio New Zealand:

- www.tvnz.co.nz/one-news/new-zealand/whitebait-eels-found-in-wellingtons-stormwater-system
- www.rnz.co.nz/national/programmes/ourchangingworld/audio/2018697287/the-streams-beneath-the-streets

6 ACKNOWLEDGEMENTS

I greatly appreciated the assistance of various WWL staff for providing early guidance and GIS information (Michelle Chew, Rhyann McCoy, Paul Davison, Jack Furnish). Megan Oliver and Evan Harrison from GWRC provided much encouragement and funded the work. Michael Greer (formerly GWRC) helped get the project off the ground in the early stages. Thanks to Renee Wilkie and the crew from Silver Linings Contracting who ensured the fieldwork was safe, efficient, and enjoyable. Thank you to my EOS Ecology colleagues for peer review (Shelley McMurtrie) and GIS assistance (Kirsty Brennan).

7 REFERENCES

Harding, J.S., Clapcott, J.E., Quinn, J.M., Hayes, J.W., Joy, M.K., Storey, R.G., Greig, H.S., Hay, J., James, T., Beech, M.A., Ozane, R., Meredith, A.S. & Boothroyd, I. 2009. *Stream Habitat Assessment Protocols for Wadeable Rivers and Streams of New Zealand*. School of Biological Sciences, University of Canterbury, Christchurch, New Zealand. 133 p.

Stark, J.D. & Maxted, J.R. 2007b. A User Guide for the Macroinvertebrate Community Index. Cawthron Institute, Nelson. Report No. 1166. 66 p.

www.eoecology.co.nz
info@eoecology.co.nz

PO Box 4262, Christchurch 8140
P 03 389 0538

PO Box 8054, Palmerston North 4446
P 06 358 9566



SCIENCE + ENGAGEMENT



Freshwater Ecology of Piped Streams in Wellington: Pilot Study Final Report

EOS Ecology Report No. GRE01-17087-02 | April 2020

Prepared for Greater Wellington Regional Council

Prepared by EOS Ecology – Dr Alex James

Reviewed by Shelley McMurtrie (EOS Ecology) and Dr Evan Harrison (GWRC)



www.eosecology.co.nz
info@eosecology.co.nz

PO Box 4262, Christchurch 8140
P 03 389 0538

PO Box 8054, Palmerston North 4446
P 06 358 9566

SCIENCE + ENGAGEMENT



EOS Ecology or any employee or sub-consultant of EOS Ecology accepts no liability with respect to this publication's use other than by the Client. This publication may not be reproduced or copied in any form without the permission of the Client.

© All photographs within this publication are copyright of EOS Ecology or the credited photographer; they may not be used without written permission.

EXECUTIVE SUMMARY	1
1 INTRODUCTION.....	2
2 METHODS.....	3
2.1 Health and Safety	3
2.2 Site Selection.....	3
2.3 Detailed Ecological Survey	6
2.4 Data Analysis.....	7
3 RESULTS.....	9
3.1 Habitat.....	9
3.2 Fish.....	12
3.3 Benthic Macroinvertebrates – Surber Sampling.....	15
3.4 Macroinvertebrates – Sticky Traps	18
3.5 Benthic Macroinvertebrates – Surber vs. Kick Net Samples	20
3.6 Benthic Macroinvertebrates – Piped vs. Open Stream Sites	24
4 CONCLUSIONS	29
4.1 Habitat.....	29
4.2 Fish Assemblages.....	31
4.3 Benthic Macroinvertebrate Assemblages	32
4.4 Sampling Methodologies.....	33
4.5 Recognition and Protection.....	35
5 RECOMMENDATIONS.....	36
5.1 Sampling Methodologies.....	36
5.2 Habitat Improvements.....	36
5.3 Recognition and Protection	37
5.4 Further Research	38
6 MEDIA.....	38
7 ACKNOWLEDGEMENTS	39
8 REFERENCES.....	39

9	APPENDICES	40
9.1	Appendix 1 – SIMPER Results Comparing Sites.....	40
9.2	Appendix 2 - BEST (Biota and Environment Matching) Results.....	43
9.3	Appendix 3 – SIMPER Results Comparing Catchments.....	44
9.4	Appendix 4 – SIMPER Results Comparing Open and Piped Stream Sites	45

EXECUTIVE SUMMARY

The piping of natural stream channels is commonplace in many urban environments. Many catchments are extensively piped such that original open stream habitat only exists as isolated remnants among the pipe network. In Wellington city, with its steep topography and small coastal catchments, many small streams have undergone extensive piping as the landscape was urbanised. Because of the difficulty of sampling piped streams, most freshwater ecological information from these urban streams is derived from remnant open sections, even though these only account for a small proportion of total stream length. To address this knowledge gap, EOS Ecology and Greater Wellington Regional Council (with support from Wellington City Council) undertook a pilot study to investigate the habitat, fish communities, and macroinvertebrate assemblages of six piped stream sites across three predominantly urban catchments in Wellington.

In-pipe habitat ranged from fast, laminar flows over bare brick or concrete, to zones of slower water velocity with natural stony substrates. Organic material such as leaves, woody debris, and tree roots were present to some extent at all sites, while biofilms were prominent at five of the six sites. Water depths were generally very shallow (<10 cm) and fish cover lacking. Just as in natural stream channels, it is apparent piped streams have erosional and depositional zones.

Fish were found at five of the six sites, indicating fish, in particular eels (both longfin and shortfin), are relatively common and widespread in piped streams. A single young adult inanga was found at a tidally influenced site in Miramar, while two banded kokopu (live juvenile; dead, recently predated adult) were found further up the same catchment. Eels were often found hiding under whatever cover they could find (i.e., rubbish, rubble, tree roots, fine sediments), indicating that even in near complete darkness they prefer to spend some of their day concealed. A comparison of the fish assemblages of piped stream sites with open stream sites in the same catchments indicated strong habitat partitioning, with eels dominating piped sites and banded kokopu dominating open sites.

The macroinvertebrate community of piped streams was dominated by taxa known to be tolerant or to prefer degraded stream conditions (e.g., *Potamopyrgus* and *Physa* snails, oligochaete worms, Orthocladinae and *Polypedilum* midge larvae, Collembola springtails, Psychodidae gnat larvae, mites, and *Ferrissia* limpets). All piped stream sites had mean MCI and QMCI values indicative of poor conditions. When compared to macroinvertebrate data collected by Greater Wellington Regional Council (GWRC) at open stream sites in the same catchments, piped stream sites had lower numbers of individuals, lower taxa richness, lower MCI and QMCI scores, and lower EPT abundance and richness. Piped stream macroinvertebrate communities were dominated by the most tolerant taxa that are capable of living in near complete darkness, with no riparian vegetation or benthic algae, and subject to a very flashy flow regime. In catchments with extensive piping, even small sections of remnant open channel are very important at maintaining catchment aquatic macroinvertebrate diversity, in particular EPT taxa.

Currently, piped streams are generally considered part of the stormwater network and are not recognised for their ecological values by the Proposed Natural Resources Plan. Should a piped stream be disturbed by some activity (e.g., construction or discharge), there is no requirement for ecological values to be taken into account or even for GWRC freshwater scientists to be notified. It is recommended that work be undertaken to allow for piped streams to be formally identified, their ecological values documented, and steps taken such that they are included in the next iteration of the Natural Resources Plan.

In-pipe habitats could be improved by the retrofit of various features such as flexible baffles in zones of laminar, high flow velocities to improve fish passage and cover for resident fish, installation of small structures to increase water depths (e.g., leaky weirs, vanes), and installation of secure artificial fish cover elements (e.g., small pipes and half pipes). Such features may also facilitate the retention of natural substrates and organic material, further improving habitat quality, including habitat for aquatic macroinvertebrates.

1 INTRODUCTION

In most cities urban streams have been extensively piped to provide land for development and alleviate flood and disease risk. In many catchments open stream habitat exists only as short remnant channels. To date ecological information on these highly modified catchments has primarily been derived from these remnant open sections for reasons of accessibility and safety, even though these sections make up a small percentage of total stream length. Further, the piped sections that join isolated open reaches are generally managed as part of the stormwater network and have typically been ignored from an ecological perspective.

Wellington, with its steep topography and coastal location, is a good example of a city where numerous small coastal stream catchments have been extensively piped, such that in several suburbs open stream channels are now only found as fragmented remnants. With the exception of some anecdotal reports of eels down stormwater grates, piped streams in Wellington have previously only been considered as migration pathways for some freshwater fishes that are known to be present in remnant open sections (e.g. banded kokopu, kōaro, eels). The fish and macroinvertebrates living within the piped sections are unknown and there has never been any attempt to characterise freshwater habitat condition in these highly modified stream environments.

EOS Ecology undertook a series of urban stream catchments investigations from 2015 to 2017 for Wellington Water Limited (WWL) to support Integrated Catchment Management Plan (ICMP) development, where the project brief was only to examine open sections of streams (James, 2016; James, 2017a; James, 2017b; James, 2018a; James, 2018b). It became apparent the ecological values of extensively piped catchments could not be fully determined without examining the piped sections (which were often the higher proportion of total stream length). Based on the catchment knowledge gained during the ICMP investigations, EOS Ecology then developed a plan to undertake a pilot study to survey piped stream ecology in Wellington. The lack of information on piped stream ecology was suggested to Greater Wellington Regional Council (GWRC) as a major knowledge gap and they agreed to fund (with assistance from Wellington City Council) a pilot study of six sites focussing on fish, macroinvertebrates, and habitat quality.

EOS Ecology was contracted by GWRC to design, implement, and report on this pilot study. Following on from an interim report completed in July 2019 (James, 2019), this final report incorporates analysis and interpretation of the quantitative Surber sampling macroinvertebrate data from the detailed survey. We examine habitat conditions, the fish community, and the macroinvertebrate assemblage. We compare piped site fish data with that from open stream sites in the same catchments. Analysis of the macroinvertebrate assemblage of the six sites based on the quantitative Surber sample information also includes comparison with data obtained earlier via modified long-handled kick nets deployed from the street surface, along with open stream standard kick net data from within the same catchments obtained from the GWRC urban stream sampling programme.

2 METHODS

2.1 Health and Safety

Underground pipes are classified as confined spaces, meaning only appropriately trained persons are able to enter and special procedures are required (e.g., operating a permitting system, use of gas detectors, use of a winch and rescue-tether system). As WWL manages the piped stream network as part of the stormwater system, a WWL-approved contractor was required to assist with the project. WWL put us in touch with Silver Linings Contracting, who were then contracted to assist with all piped stream fieldwork. They were in charge of site health and safety and supplied all equipment required for confined space entry. From EOS Ecology, the author (Alex James) was confined spaces entry certified.

2.2 Site Selection

2.2.1 Desktop Exercise

WWL provided GIS layers of the stormwater network, manhole locations, and remnant open channels. These were overlain with Google Earth imagery and used to select manholes within Wellington's inner suburbs where entry for an ecological survey appeared logistically realistic. Criteria for potential sites included:

- » Pipe diameters of no less than 1200 mm.
- » Manholes to be located on footpaths or berms to avoid the requirement for traffic management or entering of private property (Figure 1).
- » The catchment would preferably have permanently flowing open stream remnants from which fish were known.

Based on these criteria, 36 candidate manholes were identified with the expectation some would be inaccessible in the field.



Miramar Stream manhole on grassed berm



Pae Kawakawa Stream manhole on footpath

Figure 1 Examples of berm and footpath locations of candidate manholes.

2.2.2 Manhole Lifting Exercise

Over 1–2 May 2018, a manhole lifting site visit was undertaken with Silver Linings Contracting to aid selection of the final survey sites. Overall, 20 manholes were opened, with the remaining 16 either being inaccessible (e.g. buried, car parked over, building on top) or being in close proximity to another opened manhole. We also took the opportunity to trial macroinvertebrate sampling from the surface utilising a modified sampling net.

Once each manhole was opened, a set procedure was undertaken:


- » Lowering of a gas detector to check for unsafe concentrations of oxygen, hydrogen sulphide, carbon monoxide, and flammable gases (Figure 2).
- » Measurement of surface to pipe bottom and pipe diameter using a Leica Disto laser measurement device.
- » Visual estimation of water depth (dry/shallow/medium/deep), substrate type (brick, concrete, silt, cobbles, other), pipe shape/profile (circular/rectangular/arch/other), and pipe material (concrete/brick/other).
- » Collection of macroinvertebrate samples using a brush and standard kick net with extendable handles (2.7–5 m). Such samples were collected from 16 sites (Figure 2). Samples were preserved with isopropyl alcohol (IPA) for later processing.
- » The taking of photos of the manhole shaft and stream below.
- » The taking of video footage from selected manholes using a GoPro video camera and torch attached to the extendable-handle kick net (Figure 2).




Using the extendable-handle kick net to collect a macroinvertebrate sample



Using the extendable-handle kick net in conjunction with a brush to collect a macroinvertebrate sample



Measuring the atmosphere using a gas detector



The GoPro video camera setup

Figure 2 Examples of manhole lifting exercise methodology.

2.2.3 Final Site Selection

Six sites were selected for detailed ecological survey: three in Pae Kawakawa Stream catchment (Island Bay), two in Miramar Stream (Miramar), and one in Waipapa Stream (Hataitai) (Figure 3). The Pae Kawakawa Stream and Miramar Stream sites were all vertical entries down manholes, while Waipapa Stream involved entry via the ocean outlet at low tide. It was originally planned to have a site in the Waitangi Stream catchment, however the only suitable site was deemed too dangerous to enter without using breathing apparatus, for which additional specialist training would have been required. One of the three Pae Kawakawa Stream sites was a substitute for this site.



Figure 3 Locations of the six detailed survey sites and the two GWRC open stream sites in the Pae Kawakawa, Miramar and Waipapa catchments. The GWRC site codes are shown in parentheses.

2.3 Detailed Ecological Survey

The six sampling sites were visited on two occasions for the detailed survey (Stage 1: 6–7 March 2019 and Stage 2: 20–21 March 2019). Stage 1 involved installation of trail/game cameras, sticky traps, and collection of macroinvertebrate samples, while Stage 2 consisted of the recovering of cameras and sticky traps and undertaking a fish survey. In practice, fish were often sighted during Stage 1, so an informal fish survey (consisting of noting if fish were seen) was also undertaken at that time. Additionally, an equipment mishap required macroinvertebrates at one site to be collected on the Stage 2 visit. Details of the survey methodology are detailed below:

- » **Trail/game cameras:** At each site, a single trail camera was installed near the manhole, with the lens facing towards the wetted channel in an attempt to obtain images of fish within the piped streams (Figure 4). Cameras were mounted on existing pipes or reinforcing steel rods where available, otherwise a steel extendable curtain rod was affixed across the width of the piped stream. Curtain rods were attached using cable ties to any available protrusions (e.g., ladder rungs, reinforcing steels) and/or appropriate concrete adhesives applied via caulking gun. Cameras were set to take time-lapse images every 10 minutes for the duration of deployment and also be triggered to take a photo by movement. The typical movement sensor of such cameras relies on a difference in temperature between the environment and the animal, hence, is unlikely to be triggered by fish. Where the camera model allowed (we used three different models), 10-second videos were also recorded every 10 minutes.
- » **Sticky traps:** At each site, a single 24.5 cm wide by 40 cm high plastic yellow sticky trap was installed as near the roof of the piped stream as possible (Figure 4). These were sticky on both the upstream and downstream sides and installed to existing pipes or reinforcing steels at some sites or to the curtain rod described above at others.
- » **Macroinvertebrates:** Twelve Surber samples (0.09 x 0.09 m²) were collected from each site over a 40 to 100 m length of stream, depending on availability of habitat suitable for Surber sampling and access (Figure 4). Samples were preserved with isopropyl alcohol (IPA) for later processing. Macroinvertebrate samples were processed using a full-count method.
- » **Fish survey:** At each site, a 200 m (100 m upstream and downstream of the manhole access point) reach was carefully searched for fish using a spotlight and hand netting technique (Figure 4). The exception was Waipapa Stream, where we entered via the ocean outlet. At this site, we searched for fish from the entrance all the way upstream to where the macroinvertebrate survey was undertaken. In practice, in Waipapa Stream, fish were only observed in the lower 90 m of pipe as above this point water became too shallow and cover lacking for the fish present. Any obvious fish cover elements (e.g., pieces of wood, brick, and assorted rubbish) were slowly lifted to determine if fish were underneath. Fish were identified to species, with the exception of those eels that were unable to be captured. Lengths were visually estimated.
- » **Habitat characterisation:** At each Surber sample location (12 at each site), water depth, wetted width, substrate composition, and organic matter cover (leaves, CPOM, biofilms) was recorded. We recorded the “head” where water depth was recorded to enable an estimate of water velocity using the ruler methodology (cf. Harding et al., 2009 – Appendix 3). Biofilm colour and thickness was also recorded. Site photos were taken, including images of each Surber location prior to sampling (Figure 4). We also made site wide measurements of meso-habitat length (rapid, run, riffle, pool, fall) and made visual estimates of substrate composition and organic matter cover (leaves, roots, CPOM, biofilms). Any potential fish barriers were noted.
- » **Water quality:** A YSI multiprobe supplied by GWRC was used to collect spot water quality records of temperature, dissolved oxygen, and conductivity.




Camera and sticky trap attached to reinforcing steel



Surber sampling



An eel captured via spotlighting and hand netting



Example of substrate sampled for macroinvertebrates

Figure 4 Images of detailed survey methodologies.

2.4 Data Analysis

Habitat data were summarised by calculating mean values for those parameters measured at each of the 12 Surber sampling locations (water depth, wetted width, water velocity, substrate composition, and organic matter cover). Fish data were summarised by total abundance, taxa richness, the density of fish captured per linear length of pipe sampled, and relative abundance.

Macroinvertebrate data were summarised by total density, taxa richness, the densities or abundance of the five most of abundant taxa, and non-metric multidimensional scaling ordination (NMS). Biotic indices calculated were the number of Ephemeroptera-Plecoptera-Trichoptera taxa (EPT richness), % EPT abundance (i.e., EPT abundance/total abundance $\times 100$), the Macroinvertebrate Community Index (MCI), and its quantitative derivative (QMCI). The paragraphs below provide clarification on some of these metrics.

Total density refers to the number of macroinvertebrates in the sampled area for each 0.1 m² Surber sample. Taxa richness is the number of different taxa identified in each sample. 'Taxa' is generally a term for taxonomic groups, and in this case refers to the lowest level of classification that was obtained during the study. Taxa richness can be used as an indication of stream health or habitat type, where sites with greater taxa richness are usually healthier and/or have a more diverse habitat.

EPT refers to three Orders of invertebrates that are generally regarded as 'cleanwater' taxa. These Orders are Ephemeroptera (mayflies), Plecoptera (stoneflies), and Trichoptera (caddisflies), forming the acronym EPT. These taxa are relatively intolerant of organic enrichment or other pollutants and habitat degradation. The exceptions to the rule are hydroptilid caddisflies (e.g., Trichoptera: Hydroptilidae: *Oxyethira*, *Paroxyethira*), which are algal piercers and

often found in high numbers in nutrient enriched waters. Hydroptilids were not found at the piped stream sites, nor at the GWRC open stream sites. EPT richness and % EPT scores can provide a good indication as to the health of a particular site. EPT taxa are generally more diverse in non-impacted, “pristine” stream systems.

The MCI score (and its quantitative variant, QMCI) can be used to determine the level of organic enrichment for stony-bottomed waterways in New Zealand (Stark, 1985). It calculates an overall score for each sample, which is based on pollution-tolerance values for each macroinvertebrate taxon that range from 1 (very pollution tolerant) to 10 (pollution-sensitive). MCI is calculated using presence/absence data, whereas the QMCI score incorporates abundance data and so gives a more accurate result by differentiating rare taxa from abundant taxa. According to the interpretative categories of Stark & Maxted (2007), excellent conditions are indicated by a QMCI of >5.99 and an MCI of >119; good conditions by a QMCI of 5.00–5.99 and an MCI of 100–119; fair conditions by a QMCI of 4.00–4.99 and an MCI of 80–99; and poor conditions by a QMCI of <4.00 and an MCI of <80. GWRC has classified streams in their region based on their physical characteristics and developed river class-specific “MCI ecological condition classes” (Clapcott & Goodwin, 2014). The sampled streams were all classified as “Class 6 – Lowland, small”, which has MCI classes nearly identical to those of Stark & Maxted (2007) (i.e., <80 = poor, 80–100 = fair, 100–120 = good, and ≥ 120 = excellent).

NMS is non-metric statistical technique that condenses sample data (in this case macroinvertebrate community data) to a single point in low-dimensional ordination space using some measure of community dissimilarity (Bray-Curtis metric in this instance). Interpretation is straightforward such that points on an x-y plot that are close together represent samples that are more similar in community composition than those further apart (Clarke & Gorley, 2006). Significant differences in macroinvertebrate community composition between sites, catchments, Surber vs. kick net samples, and piped vs. open streams were tested using the analysis of similarities (ANOSIM) procedure, which is a non-parametric procedure, applied to the similarity matrix that underlies the NMS ordination. ANOSIM is an approximate analogue of the standard ANOVA (analysis of variance) and compares the similarity between groups (in this instance upstream control and downstream impact) using the R test statistic. $R=0$ where there is no difference in macroinvertebrate community between groups, while $R=1$ where the groups have completely different communities. Where ANOSIM results showed significant or near-significant differences in macroinvertebrate community compositions, the similarity percentages (SIMPER) procedure was used to determine which taxa were responsible. NMS, ANOSIM, and SIMPER were all carried out in PRIMER v6.1.5 (Clarke & Gorley, 2006). To determine which of the measured environmental variables (wetted width, water depth, water velocity, biofilm cover, substrate index) best explained the macroinvertebrate assemblages observed, the BEST procedure in PRIMER v6.1.5 was used (Clarke & Gorley, 2006).

3 RESULTS

3.1 Habitat

Water depths were generally very shallow, with only one site (the tidally influenced Waipapa Stream) having an average water depth greater than 0.1 m (Table 1). Two sites had relatively swift water velocities (The Parade – Dover St and Waipapa), which were a result of pipe gradient. The other sites had very low water velocity, with the 302 The Parade and Miramar – Shops sites having no detectable water velocity with the velocity head rod methodology used (Table 1). Two sites had virtually no mobile substrate, with the streambed being entirely bare concrete (The Parade – Dover St and) or bricks (Waipapa) (Table 1, Figure 5). These were also the two sites with the relatively high water velocities; hence it appears there are minimal depositional areas in these high gradient sections. The 302 The Parade site was also predominantly concrete but did have some areas of deposited substrate. The three remaining sites had a significant cover of a range of stony and sandy substrates (Table 1, Figure 5). Biofilms were prominent at all sites, with the exception of 348 The Parade where none were detected (Table 1). Other organic matter such as leaves and wood were relatively uncommon. Mesohabitat was predominantly runs, with the only riffle-type habitat present at 348 The Parade. Deeper pool habitat was only present at 302 The Parade and 348 The Parade (Table 1).

Spot water temperatures were in the 16.4–18.5 °C range across the sites, while specific conductivity was very high (28,440 $\mu\text{S}/\text{cm}$) at the Miramar – Shops site due to the influence of seawater (Table 1). The other five sites had relatively high specific conductivities (323–481 $\mu\text{S}/\text{cm}$) for freshwater systems. All sites had well oxygenated water, with only the tidally influenced Miramar – Shops site being less than 80% saturation at the time of measurement (Table 1).

Table 1 Habitat characteristics measured at each Surber sampling location (n=12 per site), mesohabitat percentages measured over the length of the macroinvertebrate sampling reaches, and water quality spot measures recorded at the entrance manholes (or start of macroinvertebrate sampling reach in Waipapa Stream).

Parameter	Pae Kawakawa Stream			Waipapa Stream	Miramar Stream	
	The Parade – Dover St	302 The Parade	348 The Parade	Waipapa*	Miramar Park	Miramar – Shops**
Mean wetted width (m)	1.17	0.63	1.14	0.52	1.25	1.38
Mean water depth (m)	0.07	0.05	0.05	0.07	0.04	0.16
Mean water velocity (m/s)	0.51	0	0.05	0.61	0.02	0
Mean substrate composition (%)	Concrete: 100	Concrete: 77 Cobble: 2 Pebble: 3 Gravel: 5 Sand: 13	Concrete: 12 Cobble: 3 Pebble: 4 Gravel: 40 Sand: 41	Brick: 100	Cobble: 5 Gravel: 13 Pebble: 80 Sand: 2	Concrete: 25 Cobble: 5 Pebble: 37 Gravel: 31 Sand: 2
Mean organic matter cover (%)	Biofilms: 100 Leaves: 0.1	Biofilms: 76 Wood: 0.2 Leaves: 0.7	Wood: 2 Leaves: 1	Biofilms: 90	Biofilms: 92.5 Wood: 0.2 Leaves: 0.6	Biofilms: 60
Mesohabitat lengths (%)	Run: 100	Run: 97 Pool: 3	Run: 51 Riffle: 44 Pool: 5	Run: 100	Run: 100	Mesohabitat varies with tide
Spot temperature	16.4 °C	18.5 °C	18.2 °C	17.4 °C	17.7 °C	17.8 °C
Spot specific conductivity	481 µS/cm	323 µS/cm	580 µS/cm	371 µS/cm	440 µS/cm	28,440 µS/cm
Spot dissolved oxygen	83.6% 8.27 mg/L	93.4% 8.86 mg/L	83.6% 7.95 mg/L	95.2% 9.22 mg/L	84.6% 8.16 mg/L	74.8% 6.49 mg/L
Spot water quality measurement time	10:30 am	4:15 pm	3:00 pm	12:20 pm	9:00 am	2:15 pm

*The Waipapa site data presented was collected in the reach where macroinvertebrates were sampled. All fish were observed in the lower, tidally influenced 90 m of pipe, which had a mostly bare concrete base.

**The Miramar – Shops site was tidally influenced, hence wetted width and depth will vary over tidal cycles.



The Parade – Dover St



302 The Parade



348 The Parade



Waipapa Stream



Miramar Park



Miramar - Shops

Figure 5 Images of each site within the macroinvertebrate sampling reach.

3.2 Fish

Fish were present at five of the six sites that underwent detailed ecological survey, with only The Parade – Dover St site not having any fish found over the 200 m survey reach (Figure 6). A total of 54 fish were found at the five piped sites. Fish abundance ranged from 6–14 individuals, with the 348 The Parade site having the highest number of fish (15 fish) and the 302 The Parade site having the least (six fish) (Figure 6). Species diversity was low with four species being found across the sites, which were (in order of abundance) longfin eel, shortfin eel, banded kokopu, and inanga (Figure 6). Longfin and shortfin eels were found at all sites where fish were found, whilst two banded kokopu (a recently dead adult and a post-whitebait juvenile) and one inanga were only found at one site each (Figure 6, Figure 7). The two kokopu were found at the Miramar Park site. One was a recently dead adult banded kokopu (150 mm long), which, based on the markings on the body, had been killed by a large eel (Figure 6). One banded kokopu post-whitebait juvenile (50 mm long) was also found at this site. The single inanga was a young adult (70 mm) found at the tidally influenced Miramar – Shops site (Figure 6, Figure 7).




Figure 6 Fish community metrics and relative abundances from the six piped stream sites surveyed on 20–21 March 2019. ‘UnIDeel’ = unidentified eels, and refers to those eels that were not able to be caught to allow for a definitive identification. Dark blue bars indicates the Pae Kawakawa catchment, grey bars Waipapa Stream, and light blue bars the Miramar catchment.



Longfin eel



Inanga young adult



Post-whitebait juvenile banded kokopu



Banded kokopu adult (dead)

Figure 7 Fish species found during surveys of the six piped stream sites on 20–21 March 2019.

Recent GWRC fish survey data were available from open stream sites in the Pae Kawakawa (two sites) and Miramar Stream (one site), allowing for comparison of fish assemblages between remnant open stream sections and piped stream sections (Figure 8). In the Pae Kawakawa catchment, no fish (or waikōura/freshwater crayfish) were found at one open and one piped survey site. At sites where fish were found, there was a distinct difference in composition between open and piped stream sections. Shortfin and longfin eel were found exclusively at the piped sites, while the one open site with fish had banded kokopu (Figure 8). In the Miramar Stream catchment, eels (shortfin and longfin) also dominated in the piped sites and longfin eels were only found at piped sites, while the single open stream site surveyed had mostly banded kokopu (Figure 8). Waikōura were only found in open stream sites in both catchments. Overall, the addition of the piped stream surveys has increased the known diversity of fish in both catchments by two species (longfin and shortfin eel for Pae Kawakawa Stream, and inanga and longfin eel for Miramar Stream).




Figure 8 Relative abundance of fish species (including waikōura) at piped and open survey sites in the Pae Kawakawa Stream and Miramar Stream catchments. Open stream sites were surveyed by GWRC and their site codes are shown in parentheses.

Eels were also observed in trail camera footage, although it was impossible to determine species (Figure 9). Full consideration of the use of trail cameras in piped streams is given in Section 4.4.



Figure 9 Trail camera infrared night vision photo from the 348 The Parade site clearly showing an eel in midframe.

3.3 Benthic Macroinvertebrates – Surber Sampling

A total of 31 invertebrate taxa were recorded from the six sites that underwent detailed ecological survey. The most diverse groups were the two-winged flies (Diptera: 10 taxa), crustaceans (Crustacea: 5 taxa), molluscs (Mollusca: 4 taxa), caddisflies (Trichoptera: 3 taxa), and polychaete worms (Polychaeta: 2 taxa). Groups represented by one taxon included springtails (Hexapoda: Collembola), beetles (Coleoptera), mites (Arachnida: Acari), and four groups collectively called worms (Nematoda, Nemertea, Oligochaeta, Platyhelminthes).

The total number of taxa captured at each site ranged from nine (302 The Parade and Miramar – Shops) to 14 (The Parade – Dover St). Taxa unique to each site ranged from one (302 The Parade) to six (The Parade – Dover St) (Figure 10). Sites were dominated by non-insect taxa with oligochaete worms and *Potamopyrgus* snails being particularly common (most abundant taxa at three and two sites respectively) (Figure 10). Other common non-insect taxa included collembola, *Physa* snails, *Ferrissia* limpets, and mites. The only relatively common insect taxa at some sites were various Diptera (true flies) taxa (Orthocladiinae and *Polypedilum* midge larvae, Psychodidae gnat larvae). The only other relatively common insect were Scirtidae beetle larvae, which were only found at the The Parade – Dover St site (Figure 10). The Trichoptera captured were single individuals of *Hydropsyche* (*Aoteapsyche*), *Oeconesus* (both at 348 The Parade), and *Polyplectropus* (at Miramar Park). Based on such rarity, it is likely these individuals had been transported downstream from open sections upstream.

NMS ordination indicated samples from some sites were far more variable than others. For example, those samples from the Miramar catchment (Miramar Park and The Parade – Dover St) clustered fairly tightly together compared to the large spread of the 302 The Parade and Waipapa Stream sites (Figure 11). ANOSIM comparing sites indicated a moderately strong (Global R=0.63) and significant (p=0.01) difference among the sampling sites (Figure 11). SIMPER results showing the main taxa responsible for differences between each site pair are shown in Appendix Table 1. Between site differences often resulted from differences in the abundance of common, widespread taxa (e.g., Oligochaeta worms and *Potamopyrgus* snails). However, on several occasions taxa that are present in one site, but absent from another contribute to differences (often Psychodidae gnat larvae, *Physa* snails, or Scirtidae beetle larvae).

The BEST procedure indicated that of the measured environmental variables, a combination of wetted width and water velocity was the most correlated with the macroinvertebrate community data, although the correlation was not strong (0.36) (see Appendix Table 2 for full BEST results).































Pae Kawakawa Stream			Waipapa Stream	Miramar Stream	
The Parade – Dover St	302 The Parade	348 The Parade	Waipapa Stream	Miramar Park	Miramar - Shops
Total taxa: 14 Unique taxa: 6	Total taxa: 9 Unique taxa: 1	Total taxa: 13 Unique taxa: 4	Total taxa: 10 Unique taxa: 3	Total taxa: 10 Unique taxa: 2	Total taxa: 9 Unique taxa: 4
 Oligochaeta worms (MCI=1; 41%)	 <i>Potamopyrgus</i> snails (MCI=4; 46%)	 <i>Potamopyrgus</i> snails (MCI=4; 77%)	 Collembola springtails (MCI=6; 43%)	 Oligochaeta worms (MCI=1; 49%)	 Oligochaeta worms (MCI=1; 48%)
 Psychodidae gnat larvae (MCI=1; 18%)	 <i>Physa</i> snails (MCI=3; 28%)	 Oligochaeta worms (MCI=1; 15%)	 Oligochaeta worms (MCI=1; 23%)	 <i>Potamopyrgus</i> snails (MCI=4; 37%)	 <i>Polypedilum</i> midge larvae (MCI=3; 32%)
 Orthoclaadiinae midge larvae (MCI=2; 14%)	 Oligochaeta worms (MCI=1; 18%)	 <i>Physa</i> snails (MCI=3; 4%)	 Orthoclaadiinae midge larvae (MCI=2; 9%)	 <i>Physa</i> snails (MCI=3; 7%)	 Acarina mites (MCI=5; 7%)
 Collembola springtails (MCI=6; 14%)	 <i>Ferrissia</i> limpets (MCI=3; 3%)	 <i>Ferrissia</i> limpets (MCI=3; 2%)	 Psychodidae gnat larvae (MCI=1; 9%)	 <i>Ferrissia</i> limpets (MCI=3; 3%)	 <i>Potamopyrgus</i> snails (MCI=4; 6%)
 Scirtidae beetle larvae (MCI=8; 5%)	 Acarina mites (MCI=5; 2%)	 Orthoclaadiinae midge larvae (MCI=2; 1%)	 Acarina mites (MCI=5; 3%)	 Empididae fly larvae (MCI=3; 1%)	 Platyhelminthes flatworms (MCI=3; 2%)

Figure 10 The total number of taxa captured at each site, the number of taxa that were unique to that site, and the five most abundant taxa at each of the six piped stream sampling sites. The relative abundance percentages and MCI-hb scores for each taxon are shown in parentheses. All images © EOS Ecology except *Ferrissia* (A. Mrkvicka), *Polypedilum* (Landcare Research), Psychodidae (MAF), and Scirtidae (Landcare Research).

When looking at the catchment scale, there is wide scatter among the samples with the ANOSIM comparing catchments, indicating a relatively weak (Global $R=0.19$) but significant difference (Figure 12). SIMPER results showing the main taxa responsible for differences between each catchment pair are displayed in Appendix Table 3. In terms of catchment pairs, the Miramar and Waipapa catchments were the most different ($R=0.73$), while the Pae Kawakawa and Miramar catchments were the least different ($R=0.05$), with the lack of Psychodidae gnat larvae in the Miramar samples being the most notable faunal difference.




Figure 11 Non-metric multidimensional scaling (NMS) ordination of macroinvertebrate community data from the six piped stream sites with data divided by site. Each point represents a single Surber sample. The analysis was based on abundance data. Also shown is the ANOSIM site comparison result. A stress value of 0.11 is indicative of a fair ordination that can still correspond to a usable picture.




Figure 12 Non-metric multidimensional scaling (NMS) ordination of macroinvertebrate community data from the six piped stream sites with data divided by catchment. Each point represents a single Surber sample. The analysis was based on abundance data. Also shown is the ANOSIM catchment comparison result. A stress value of 0.11 is indicative of a fair ordination that can still correspond to a usable picture.

Mean macroinvertebrate densities were much greater at the 348 The Parade site than the other five sites (Figure 13). In general, macroinvertebrates were at very low densities in piped streams and Surber samples were sometimes lacking any animals at all (Waipapa Stream: 4 samples; 302 The Parade and Miramar - Shops: 1 sample each). The Waipapa Stream site had particularly low densities and was also the site with the highest water velocities and only site with a 100% brick bottom (Table 1, Figure 5). Mean taxa richness was greatest at the The Parade – Dover St site and least at the Waipapa Stream site (Figure 13). The Parade – Dover St also had the highest number of overall taxa captured (14), while 302 The Parade and Miramar - Shops had the least (nine taxa each) (Figure 10). Mean MCI tended to be slightly higher at the three Pae Kawakawa catchment sites (i.e., The Parade sites) but was well within the “poor” category (Figure 13). Likewise, QMCI scores all indicated “poor” conditions (Figure 13).




Figure 13 Mean macroinvertebrate community metrics from each of six piped streams sites. Twelve Surber samples were collected at each site. The interpretive categories of Clapcott & Goodwin (2014) and Stark & Maxted (2007) are shown on the MCI and QMCI graphs, respectively.

3.4 Macroinvertebrates – Sticky Traps

Several invertebrate taxa were captured on sticky traps hanging vertically at each piped stream site. The most common and abundant taxon was the fungus gnat family, Mycetophilidae, which were present at all sites, and do not have aquatic larvae (Table 2). They are commonly found in damp habitats where their preferred host fungi are abundant. Hence the dark, damp environment of piped streams likely suits the particular Mycetophilidae genera found. Other terrestrial taxa found included mites, beetles, and ants (Table 2). Along with the various spiders observed in the pipes, these indicate there is likely a significant terrestrial arthropod community living within piped streams.

Adults of aquatic taxa included mosquitos (four sites), non-biting Chironomids (two sites), and crane flies (one site) (Table 2). Larval mosquitos were found in Surber samples at one of the four sites where adult mosquitos were found indicating, the value of sticky trapping at detecting additional aquatic taxa that may be missed by benthic sampling. Chironomidae larvae were present at both the sites adult chironomids were found at (Table 2). These results confirm there are various insect taxa flying around in permanent darkness and terrestrial and aquatic taxa are present.

Table 2 Invertebrates captured on single, double-sided sticky traps installed at each piped stream site for an approximately two week period. The sticky trap at the Waipapa Stream site was lost. For those taxa with aquatic larvae, the presence of potential larvae in Surber samples is indicated by "(larvae present)" in the Larval Habitat column.

Site name	Taxa name	Common name	Sticky trap count	Larval Habitat
The Parade - Dover St	Diptera: Mycetophilidae	Fungus gnat	5	Terrestrial
	Diptera: Culicidae	Mosquito	2	Aquatic (larvae present)
302 The Parade	Diptera: Mycetophilidae	Fungus gnat	2	Terrestrial
	Diptera: - Sciaridae	Fungus gnat	1	Terrestrial
348 The Parade	Diptera: Chironomidae	Non-biting midges	1	Aquatic (larvae present)
	Diptera: Mycetophilidae	Fungus gnat	12	Terrestrial
	Diptera: Culicidae	Mosquito	1	Aquatic
Miramar Park	Acari	Mites	6	Terrestrial
	Diptera: Mycetophilidae	Fungus gnat	17	Terrestrial
	Coleoptera	Beetles	2	Terrestrial
	Formicidae	Ants	2	Terrestrial
	Diptera: Tipulidae	Crane flies	1	Possibly Aquatic
	Diptera: Culicidae	Mosquito	3	Aquatic
	Diptera: Chironomidae	Non-biting midges	1	Aquatic (larvae present)
Miramar - Shops	Diptera: Mycetophilidae	Fungus gnat	451	Terrestrial
	Diptera: Culicidae	Mosquito	3	Aquatic

3.5 Benthic Macroinvertebrates – Surber vs. Kick Net Samples

Surber sampling involved much more sampling effort with a greater area of habitat sampled compared to kick nets collected from the surface using the long-handled kick net illustrated in Figure 2. This resulted in Surber samples collecting higher numbers and a greater diversity of benthic macroinvertebrates than the kick nets (Figure 14, Figure 15, Figure 16). Consequently Surber sample data had between three and ten extra taxa per site that were not encountered by the single kick net sample (Figure 14, Figure 15). However, at three sites the kick net samples also contained two or three taxa that were not captured in any of the 12 Surber samples collected at each site (Figure 14, Figure 15). NMS ordination of presence-absence data showed the two methods aligned more closely at some sites than others. For example, Surber and kick net data plotted relatively close together for the 348 The Parade and Miramar – Shops sites, compared to the other two sites (Figure 17). ANOSIM comparing the two methods showed no significant difference.

Despite the disparity in sampling effort, the five most common taxa at each site were generally similar between Surber and kick net sampling. For the 348 The Parade site the four most abundant taxa, and at the Miramar Park site the two most abundant taxa, were the same for both sampling methods (Figure 14, Figure 15). Each sampling method was undertaken at a different time (Surber's: March 2019; Kick nets: May 2018), which may have played some role in the macroinvertebrates captured. However, piped streams are a more stable environment compared to open streams in terms of seasonal variables like temperature and daylight, so any seasonal variation in macroinvertebrate communities is likely to be minimal.

Overall, site MCI and QMCI scores based on a single sample (kick nets) or twelve pooled samples (Surbers) varied between the sampling methods. MCI scores for pooled Surbers were higher than those of kick net samples at three of the four sites, while pooled Surber QMCI was higher than kick net scores for two of the four sites (Figure 16). However, all scores remained well in the "poor" category, with the exception of the Surber data at the 348 The Parade site, which was slightly above the "poor"-"fair" boundary (Figure 16).





















The Parade – Dover St Surber	The Parade – Dover St Kick	348 The Parade Surber	348 The Parade Kick
Total taxa: 14 Unique taxa: 10	Total taxa: 7 Unique taxa: 3	Total taxa: 13 Unique taxa: 6	Total taxa: 10 Unique taxa: 3
 Oligochaeta worms (MCI=1; 41%)	 <i>Potamopyrgus</i> snails (MCI=4; 29%)	 <i>Potamopyrgus</i> snails (MCI=4; 77%)	 <i>Potamopyrgus</i> snails (MCI=4; 35%)
 Psychodidae gnat larvae (MCI=1; 18%)	 Oligochaeta worms (MCI=1; 24%)	 Oligochaeta worms (MCI=1; 15%)	 Oligochaeta worms (MCI=1; 26%)
 Orthoclaadiinae midge larvae (MCI=2; 14%)	 <i>Polypedilum</i> midge larvae (MCI=3; 18%)	 <i>Physa</i> snails (MCI=3; 4%)	 <i>Physa</i> snails (MCI=3; 15%)
 Collembola springtails (MCI=6; 14%)	 <i>Physa</i> snails (MCI=3; 12%)	 <i>Ferrissia</i> limpets (MCI=3; 2%)	 <i>Ferrissia</i> limpets (MCI=3; 7%)
 Scirtidae beetle larvae (MCI=8; 5%)	 Talitridae amphipods (MCI=5; 6%)	 Orthoclaadiinae midge larvae (MCI=2; 1%)	 Collembola springtails (MCI=6; 6%)

Figure 14 Pae Kawakawa Stream catchment: The total number of taxa captured at each site with each method (Surbers and kick net), the number of taxa that were unique to each method at that site, and the five most abundant taxa of each method and site. The relative abundance percentages and MCI-hb scores for each taxon are shown in parentheses. All images © EOS Ecology except *Ferrissia* (A. Mrkvicka), Psychodidae (MAF), and Scirtidae (Landcare Research).





















Miramar Park Surber	Miramar Park Kick	Miramar – Shops Surber	Miramar – Shops Kick
Total taxa: 10 Unique taxa: 6	Total taxa: 6 Unique taxa: 2	Total taxa: 9 Unique taxa: 3	Total taxa: 6 Unique taxa: 0
 Oligochaeta worms (MCI=1; 49%)	 Oligochaeta worms (MCI=1; 68%)	 Oligochaeta worms (MCI=1; 48%)	 <i>Polypedilum</i> midge larvae (MCI=3; 59%)
 <i>Potamopyrgus</i> snails (MCI=4; 37%)	 <i>Potamopyrgus</i> snails (MCI=4; 19%)	 <i>Polypedilum</i> midge larvae (MCI=3; 32%)	 Acarina mites (MCI=5; 9%)
 <i>Physa</i> snails (MCI=3; 7%)	 Collembola springtails (MCI=6; 8%)	 Acarina mites (MCI=5; 7%)	 Orthoclaadiinae midge larvae (MCI=2; 9%)
 <i>Ferrissia</i> limpets (MCI=3; 3%)	 Orthoclaadiinae midge larvae (MCI=2; 2%)	 <i>Potamopyrgus</i> snails (MCI=4; 6%)	 Oligochaeta worms (MCI=1; 9%)
 Empididae fly larvae (MCI=3; 1%)	 <i>Polypedilum</i> midge larvae (MCI=3; 2%)	 Platyhelminthes flatworms (MCI=3; 2%)	 Platyhelminthes flatworms (MCI=3; 9%)

Figure 15 Miramar Stream catchment: The total number of taxa captured at each site with each method (Surbers and kick net), the number of taxa that were unique to each method at that site, and the five most abundant taxa of each method and site. The relative abundance percentages and MCI-hb scores for each taxon are shown in parentheses. All images © EOS Ecology except *Ferrissia* (A. Mrkvicka), *Polypedilum*, and Scirtidae (Landcare Research).




Figure 16 Mean macroinvertebrate community metrics from the four piped streams sites where Surber and kick net data was available. Stripped bars are Surber samples and plain bars are kick net samples. Data from the twelve Surber samples have been pooled. The interpretive categories of Clapcott & Goodwin (2014) and Stark & Maxted (2007) are shown on the MCI and QMCI graphs, respectively.




Figure 17 Non-metric multidimensional scaling (NMS) ordination of macroinvertebrate community data from the four piped stream sites where Surber (triangles) and kick net (circles) data was available. Kick nets were single samples whilst the 12 Surber samples per site were pooled to one sample per site, and all data transformed into presence-absence. Also shown is the ANOSIM method (Surber or kick net) comparison result. A stress value of 0.04 is indicative of an excellent representation of data with no prospect of misinterpretation.

3.6 Benthic Macroinvertebrates – Piped vs. Open Stream Sites

In the Pae Kawakawa (Island Bay) and Miramar catchments, GWRC had undertaken macroinvertebrate sampling from sections of remnant open stream in January 2019, with which we could directly compare the piped stream site data. It was considered the collection of twelve quantitative Surber samples per site would provide a better representation of the benthic macroinvertebrate community of piped stream sites than would the data obtained from a single, kick net sample collected down the manhole shaft with a modified, long-handled sampling net. Hence, we have used this piped stream Surber data for this piped vs. open stream comparison. GWRC sampled one open site in each catchment (Figure 3). A second open stream site in the Pae Kawakawa Stream catchment was found to be dry when visited. In the Pae Kawakawa Stream catchment a total of 22 taxa were found across three piped stream sites (7–14 taxa range at individual sites), while 29 taxa were found at the GWRC open stream site. In the Miramar catchment and total of 19 taxa were found at the two piped stream sites (nine or ten taxa at each site), while 26 taxa were found at the GWRC open stream site.

The main differences between open and piped sites in the Pae Kawakawa catchment in terms of the most abundant taxa were the dominance of chironomid midge larvae (*Polypedilum* and *Orthoclaadiinae*) and presence of relatively pollution-sensitive *Hydropsyche* (*Orthopsycha*) larvae at the open stream site (Figure 18). At the piped stream sites, while midge larvae were also among the more common taxa, oligochaete worms and snails (*Potamopyrgus* and *Physa*) tended to be the most abundant taxa (Figure 18). While also dominated numerically by *Potamopyrgus* snails, the macroinvertebrate community of the open stream site in the Miramar catchment differed from those of the two piped sites by having two pollution-sensitive taxa (*Hydropsyche* (*Orthopsycha*) caddisflies and *Zephlebia* mayflies) as the second and third most abundant taxa (Figure 19). The sampling of piped streams sites detected an additional ten taxa

in the Pae Kawakawa catchment and an additional seven taxa in the Miramar catchment to the open stream site samples taken in each catchment.




















Pae Kawakawa – Open (Island Bay Stream trib @ Farnham Street; PA01)	The Parade – Dover St	302 The Parade Surber	348 The Parade
Total taxa: 29 Unique taxa: 16	Total taxa: 14 Unique taxa: 7	Total taxa: 7 Unique taxa: 1	Total taxa: 13 Unique taxa: 2
 <i>Polypedilum</i> midge larvae (MCI=3; 25%)	 Oligochaeta worms (MCI=1; 41%)	 <i>Potamopyrgus</i> snails (MCI=4; 46%)	 <i>Potamopyrgus</i> snails (MCI=4; 77%)
 Orthocladiinae midge larvae (MCI=2; 24%)	 Psychodidae gnat larvae (MCI=1; 18%)	 <i>Physa</i> snails (MCI=3; 28%)	 Oligochaeta worms (MCI=1; 15%)
 <i>Hydropsyche</i> (<i>Orthopsyche</i>) caddisflies (MCI=9; 8%)	 Orthocladiinae midge larvae (MCI=2; 14%)	 Oligochaeta worms (MCI=1; 18%)	 <i>Physa</i> snails (MCI=3; 4%)
 Oligochaeta worms (MCI=1; 8%)	 Collembola springtails (MCI=6; 14%)	 <i>Ferrissia</i> limpets (MCI=3; 3%)	 <i>Ferrissia</i> limpets (MCI=3; 2%)
 <i>Hydropsyche</i> (early instar*) caddisflies (8%)	 Scirtidae beetle larvae (MCI=8; 5%)	 Acarina mites (MCI=5; 2%)	 Orthocladiinae midge larvae (MCI=2; 1%)

Figure 18 Pae Kawakawa Stream catchment: The number of unique taxa at each site and the five most abundant taxa of each piped stream site and the single GWRC open stream site. The relative abundance percentages and MCI-hb scores for each taxon are shown in parentheses. All images © EOS Ecology except *Ferrissia* (A. Mrkvicka), Psychodidae (MAF), *Polypedilum* and Scirtidae (both Landcare Research).
















Miramar – Open (Maupuia Stream; MR01)	Miramar Park	Miramar – Shops
Total taxa: 26 Unique taxa: 16	Total taxa: 10 Unique taxa: 3	Total taxa: 9 Unique taxa: 4
 <i>Potamopyrgus</i> snails (MCI=4; 38%)	 Oligochaeta worms (MCI=1; 49%)	 Oligochaeta worms (MCI=1; 48%)
 <i>Hydropsyche</i> (<i>Orthopsyche</i>) caddisflies (MCI=9; 17%)	 <i>Potamopyrgus</i> snails (MCI=4; 37%)	 <i>Polypedilum</i> midge larvae (MCI=3; 32%)
 <i>Zephlebia</i> mayflies (MCI=7; 8%)	 <i>Physa</i> snails (MCI=3; 7%)	 Acarina mites (MCI=5; 7%)
 Talitridae amphipods (MCI=5; 5%)	 <i>Ferrissia</i> limpets (MCI=3; 3%)	 <i>Potamopyrgus</i> snails (MCI=4; 6%)
 <i>Polypedilum</i> midge larvae (MCI=3; 5%)	 Empididae fly larvae (MCI=3; 1%)	 Platyhelminthes flatworms (MCI=3; 2%)

Figure 19 Miramar Stream catchment: The number of unique taxa at each site and the five most abundant taxa of each piped stream site and the single GWRC open stream site. The relative abundance percentages and MCI-hb scores for each taxon are shown in parentheses. All images © EOS Ecology except *Ferrissia* (A. Mrkvicka) and *Polypedilum*, (Landcare Research).

Macroinvertebrate community metrics were all greater at the open stream sites than the piped stream sites in the Pae Kawakawa and Miramar catchments (Figure 20). While different sampling methods were used (single composite kick net at open sites; 12 Surber samples at piped sites), it is clear there were much higher numbers of macroinvertebrates and taxa at the open stream sites in both catchments. MCI and QMCI were higher at open sites, although only marginally for QMCI in the Pae Kawakawa catchment (Figure 20). EPT taxa richness and percentage EPT individuals were substantially higher at the open sites, with three of the five piped stream sites not having any EPT taxa at all (Figure 20). At those piped sites with EPT taxa, these were just single individuals that may well have been transported from upstream open sites during floods, rather than being permanent residents.




Figure 20 Macroinvertebrate community metrics from open and piped stream sites in the Pae Kawakawa (dark blue bars) and Miramar (light blue bars) catchments. Open stream data were derived from a single composite kick net sample and piped stream data from pooling 12 Surber samples. Striped bars indicate open stream sites. The interpretive categories of Clapcott & Goodwin (2014) and Stark & Maxted (2007) are shown on the MCI and QMCI graphs, respectively.

NMS ordination showed open stream sites were clearly separated from piped sites along Axis 2, although all sites (both open and piped sites) were quite separated from one another along Axis 1 (Figure 21). ANOSIM indicated a marginally significant difference between open and piped sites ($p=0.048$), while a R value of 0.56 is indicative of moderate differences in macroinvertebrate composition among site type (open and piped) (Figure 21). SIMPER results indicating the taxa most responsible for this open-piped difference are shown in Appendix Table 4. Non-insect taxa (oligochaete worms, *Potamopyrgus* snails, and *Physa* snails) tended to have higher relative abundances at piped sites, while insect taxa (*Polypedilum* and Orthocladiinae midge larvae and *Hydropsyche* (*Orthopsyche*) caddisfly larvae) had higher relative abundances at the open stream site (Appendix Table 4). Given the analysis is based on relative abundances, it is important to remember that at the open stream sites benthic macroinvertebrates were generally more abundant overall (Figure 20).




Figure 21 Non-metric multidimensional scaling (NMS) ordination of macroinvertebrate community data from open and piped stream sites in the Pae Kawakawa (dark blue symbols) and Miramar (light blue symbols) catchments. Open stream data is derived from a single composite kick net sample and piped stream data from pooling 12 Surber samples. Data was converted to relative abundance prior to analysis. Also shown is the ANOSIM open vs. piped comparison result. A stress value of 0.04 is indicative of an excellent representation of data with no prospect of misinterpretation.

4 CONCLUSIONS

4.1 Habitat

Zones of cobble and gravel substrate are present within piped streams and provide a stream bed in such areas that at least appears natural, although it will usually be underlain by concrete or sealed brick. Organic material (leaves, woody debris, tree roots) were also present. Stony substrates, leaves, and wood debris have presumably been transported from upstream remnant open stream sections and stormwater inputs. Some sites had minimal or no loose rocky substrates. It is likely that piped streams have depositional and erosional zones based on gradient and water velocity as in natural stream channels, with the difference that erosional zones will generally be bare concrete or whatever material the pipe is constructed from. Water depths at the surveyed sites were generally very shallow (<10 cm) and deeper pool habitat appeared to be very limited or absent. Brown-orange biofilms were prominent at five of the six sites and presumably are a food source for some macroinvertebrates. Spot measures of dissolved oxygen indicated relatively well oxygenated water.

From an ecological perspective, piped stream habitat could be improved from measures that increase water depths, increase the abundance of pool habitat, slow water velocities, trap natural substrates and organic material, and provide cover for fish. This is especially the case in higher gradient pipes such as observed at the Waipapa Stream and The Parade – Dover St sites, where long reaches of continuous high velocities over bare concrete or brick meant there was very limited habitat for fish and invertebrates (Figure 22). Proven technologies such as flexible plastic baffles are available to retrofit such zones to improve fish passage and provide slower velocity habitat for resident fish. Such baffles may also facilitate the retention of natural substrates and organic material, further improving habitat quality, including habitat for aquatic macroinvertebrates.

In terms of deeper pool habitat, more substantial structures such as leaky rock weirs could be used in appropriate locations to create such habitat in existing pipes. Some manholes, such as that at the Miramar Park site, were designed such that below the manhole was a pool-like zone of deeper water compared to the habitat directly upstream and downstream (Figure 23). There is the potential where pipes are being replaced or new ones laid that special pipe sections that create deeper pool habitat could be incorporated into the design.



Figure 22 Long, homogenous zones of high water velocities at the Waipapa Stream site (left) and The Parade – Dover St site in the Pae Kawakawa catchment (right).




Figure 23 Pipe design at the Miramar Park site allowed the development of deeper pool-like habitat.

4.2 Fish Assemblages

Fish were present at five of the six detailed survey sites, hence appear to be widespread in the piped stream network. Fish diversity of piped stream sites was low and dominated by eels, with both longfin and shortfin eels present. Inanga appear to have the ability to live within piped streams. However, in catchments that are piped to the ocean it is highly unlikely they could ever successfully spawn due to a lack of spawning habitat. Additionally, it is highly likely they would be at low densities, would have limited ability to get beyond velocity and physical barriers due being relatively weak swimmers, and at high risk of eel predation given the lack of cover. There appears to be clear habitat partitioning of the fish fauna (and waikōura) in catchments that have extensive piped sections and remnant open headwater sections. Eels dominate the pipes while banded kokopu (and also waikōura) dominate the open sections, hence even relatively short sections of remnant open streams are crucial to maintaining catchment fish diversity.

Based on previous open stream fish survey data, it is clear the whitebait of at least banded kokopu and kōaro travel through the piped stream network to reach open sections with suitable habitat. Eels resident in the pipes likely feed extensively on upstream migrating whitebait and elvers at certain times of the year. The piped stream sites generally lacked fish cover and it was observed that even in near complete darkness, eels preferred to spend some of their time hiding under whatever cover was available, including tree roots, rubbish, concrete rubble, fine sediment accumulations, and woody debris (Figure 24). There is the potential that habitat improvement features could be added to piped streams to both provide cover for resident fish and improve conditions for migrating whitebait (e.g. fitting flexible baffles to sections with high velocity, laminar flows; increasing water depths/creating pools using leaky weirs; installing artificial fish cover elements such as securely attached small lengths of pipe).



Figure 24 Eel attempting to use tree roots as cover.

4.3 Benthic Macroinvertebrate Assemblages

Piped stream sites were dominated by oligochaete worms, snails (*Potamopyrgus* and *Physa*), and midge larvae (Orthocladiinae and *Polypedilum*). Collembola springtails, Psychodidae gnat larvae, mites, and *Ferrissia* limpets were prominent at some sites. These are taxa known to be tolerant of or that prefer degraded stream habitats, hence it is no surprise they are able to persist in piped streams. It is notable that some of these taxa typically graze on algae or plant material in open streams (e.g. snails and limpets), hence they must be able to survive on the biofilms that grow in near complete darkness and/or organic material (e.g. leaves) that enter and are retained in the piped streams. There were some between-site differences in macroinvertebrate communities, often driven by differences in the abundance of the most common and widespread taxa (e.g. oligochaete worms and *Potamopyrgus* snails). However, on several occasions taxa that were present in one site, but absent from another, contributed to differences (often Psychodidae gnat larvae, *Physa* snails, or Scirtidae beetle larvae). Of the measured environmental variables, a combination of wetted width and water velocity showed the strongest correlation with macroinvertebrate data, although this was fairly weak (0.36). Water velocity appeared to have some effect on the community composition, with the two sites with the highest water velocities and a general lack of natural substrates (Waipapa Stream and The Parade – Dover St) being the only sites not to have *Potamopyrgus* snails among the five most abundant taxa. Additionally, both these sites were also the only ones to have collembola springtails, Psychodidae gnat larvae, and Orthocladiinae midge larvae among the top five most abundant taxa. The only other known published aquatic macroinvertebrate data from New Zealand piped streams are that of Neale & Moffett (2016), who collected samples from two piped sites as they were being removed as part of a daylighting project in Auckland. One of their sites was strongly dominated by oligochaete worms followed by mollusca and Diptera larvae, while the other was dominated by Diptera larvae. Three of our six sites were also dominated by oligochaete worms, however none were by Diptera larvae, although they were present at all sites. Neale & Moffett's (2016) study catchment had far more open channel length remaining than the catchments of this pilot study, and this may have some effect on the macroinvertebrate assemblages they encountered.

Mean invertebrate densities and taxa richness were low at all piped stream sites and MCI and QMCI were well within the "poor" quality class, although it is worth noting MCI was never intended for use in artificially piped streams, hence scores may not be overly applicable to such environments. It is clear that extensive piping of the stream channel has resulted in a macroinvertebrate community being limited to those species able to persist in a totally enclosed stream with near total darkness, no riparian vegetation or benthic algae, and subject to very flashy flow conditions. Additionally, given the predominantly urban catchment, they are further subjected to the various contaminants that are common with such a land use (e.g. heavy metals, hydrocarbons, detergents, paint wash water, etc.).

A comparison of macroinvertebrate communities between open and piped stream sites in the same catchments indicated open stream sites have a greater abundance of macroinvertebrates, greater taxa richness, and more pollution-sensitive EPT taxa than piped sites. It is likely the more natural conditions of the open sites (e.g. riparian vegetation, benthic algae, day-night cycle, greater instream habitat variability and likely a more stable habitat under flood conditions) supports higher densities and a greater diversity of macroinvertebrate taxa. Longitudinal studies centred on natural cave stream entrances have shown similar results with lower invertebrate richness and diversity within cave streams (e.g. Watson, 2010). The main macroinvertebrate community differences between open and piped stream sites in the Pae Kawakawa catchment were *Polypedilum* and Orthocladiinae midge larvae being the two most abundant taxa and the relatively pollution-sensitive *Hydropsyche* (*Orthopsyche*) being the third most abundant taxa at the open site. Similarly, the main macroinvertebrate community differences in the Miramar catchment are the presence of *Hydropsyche* (*Orthopsyche*) caddisfly larvae and *Zephlebia* mayfly larvae as the second and third most abundant taxa at the open site. The open-piped site comparison in the Miramar catchment is confounded somewhat as the open site was in a headwater stream in a vegetated reserve with minimal urban inputs, while the piped sites were further downstream and subject to substantial urban stormwater inputs. This was not the case in the Pae Kawakawa catchment as the open site was in a short section with pipes upstream and downstream and thus received

urban stormwater runoff. Based on this limited data, it would appear that within extensively piped catchments, remnant sections of open stream are extremely important in maintaining catchment aquatic macroinvertebrate diversity. As such, the piping of even short sections of remnant open stream in urban catchments should be resisted. Additionally, it is evident the daylighting of piped streams, even relatively short sections, could have positive effects on catchment macroinvertebrate diversity.

For short sections of remnant open stream that have piped sections upstream and downstream (such as the GWRC Pae Kawakawa Stream site), the source of aquatic insects with flying adults (e.g., *Hydropsyche* (*Orthopsyche*)) pose several questions. Do adults fly up or down pipes to reach the site? Do adults fly across the heavily urbanised catchment to reach the site? Is the population sustained predominantly by emergence and oviposition of local individuals? Are some larvae transported to the site through the pipe network from upstream open stream habitat? Additionally, are some macroinvertebrate taxa that prefer open stream habitat able to persist in the pipes for some distance directly downstream of remnant open sections? This final question could be addressed by finding accessible pipe sites with open habitat directly upstream so you could collect samples from open habitats and at increasing distances downstream in the pipe network, utilising a combination of benthic and drift sampling methodologies.

4.4 Sampling Methodologies

For future piped stream macroinvertebrate surveys, the selection of methods would primarily depend on the overall aim of the study. Collecting a kick net sample from the surface is relatively quick and does not require confined space entry so many sites could theoretically be done in a single day. We have shown such a method can also yield macroinvertebrate community data at some sites that is fairly similar to that collected by Surber samples, at least in terms of presence/absence of more abundant taxa. There was some within-site variation in MCI and QMCI values between the two sampling techniques, although these were not great enough to change quality class interpretation with all sites being “poor”, with the exception of MCI at one site (348 The Parade) where the Surber MCI was just above the “fair” threshold. Additionally, surface kick net sampling can be used at sites that are either too small or deemed too hazardous for people to enter (i.e., smaller dimension pipes, deep pipes, pipes with inflows from former landfill sites). Further, improvements to sampling equipment, such as a curved net frame for round pipe sites and a long handled stiff brush to scrub the stream bed, could improve sampling efficiency. However, the efficiency of sampling will never match that of standard open stream kick net sampling, such that a single manhole site sampled from the surface is not directly comparable to a single open stream site. To characterise the piped stream macroinvertebrate assemblage of a catchment surface kick net samples from several separate manholes would be required.

If a more detailed and accurate representative sample of the macroinvertebrate community is desired, then full pipe entry to undertake quantitative Surber sampling is recommended. Such sampling is less awkward, enables a larger area of stream bed and differing habitats/substrates to be sampled, but it is more time consuming/expensive. However, it is evident piped stream macroinvertebrates are very sparse, so it is recommended a greater area of stream bed is sampled than this pilot study for any future in-pipe Surber sampling. Twelve samples would still be reasonable, but samples could be obtained by pooling four standard Surber areas (i.e., sample four locations for each sample), which would result in four times the area being sampled as the current study. In-pipe Surber sampling also allows direct access to measure habitat variables and undertake spotlighting fish surveys.

The use of trail cameras to detect fish was trialled. While it did not appear eel movement was sufficient to trigger their operation, footage obtained through time lapse still and video imagery clearly was able to show eels moving around in the piped stream, although the species could not be identified (Figure 9). Hence, well installed trail cameras could be used to detect the presence of eels in piped streams, and could be especially useful in areas where the pipes are too small to be entered. It would be key to install cameras in such a way that they are well above any likely high flow events and positioned to provide the best footage possible as the confined space of piped streams can result in overexposed

imagery if the infrared flash is reflected off the walls. There is the potential some kind of rig could be constructed so that cameras could be installed without the need for confined space entry. However, the use of cameras will not be sufficient to identify the species of eel, and nor would it be sufficient to see other fish, which would be too small and/or too cryptic to be seen from the cameras. Concomitantly, observations down manhole shafts without entering the piped streams can detect the presence of larger eels if they happen to be present at that particular time in the small section of stream that is visible, but are unlikely to reliably determine the species of eel or detect smaller and more cryptic fish species (e.g., banded kokopu, inanga). Thus if the aim of the study was to identify/detect all fish that may be found in a piped network, then accessing the stream to walk the piped channel would be required.

Given that the greatest expense in any pipe survey is the entering of the pipe (as this requires more personnel and specialist training and certification), then if a pipe was being entered to survey fish, then it would make sense to also collect invertebrate samples at the same time, or vice versa.

The sticky traps trialled captured relatively small numbers of flying insects, with taxa with terrestrial larvae being the most commonly trapped taxa at all sites. Despite this, they did capture taxa with an aquatic larval stage that were additional to those captured in benthic Surber samples (i.e., adult mosquitoes), although these flying adults did not necessarily originate from the surveyed section of piped stream. If sticky trapping was to be used it is recommended that several replicate traps are set along a 200 m survey reach. This would, however, require return entry to the sites after some period of time to retrieve the traps and would not be worthwhile if single-visit fish and macroinvertebrate sampling visits were being undertaken, given the expense of piped stream entry. It could be possible to design an active light trapping setup that could be suspended from manholes and thus not require piped stream entry to install and retrieve. Such a setup would require design and testing.

4.5 Recognition and Protection

There are various stormwater pipe management and construction practices that could have negative impacts on the biota that exist there:

- » The removal of accumulated sediment and detritus via suction truck will remove habitat and likely lead to fish and macroinvertebrate mortality.
- » Pipe repairs or alterations using mortars and grouts have the potential to negatively impact water quality and harm biota.
- » Locations where pipe diameter changes have the potential to create fish barriers, as observed in Pae Kawakawa Stream under The Parade in Island Bay (Figure 25).
- » The use of smooth, circular pipes may reduce the accumulation of natural gravels and detritus that provide habitat and cover for piped stream biota compared to older box culvert designs.

The permanent piping of existing remnant open stream sections will permanently alter the ecological function and fauna of those sections. There can be pressure to allow piping of such sections given that the majority of stream length in that catchment are already piped. However, this should be resisted from an ecological perspective given that these open remnant sections represent biodiversity 'hotspots' compared to piped sections for fish, macroinvertebrates, and likely algal species.

Currently, piped streams in Wellington are generally considered part of the stormwater system and managed as such. The Proposed Natural Resources Plan and the previous Regional Freshwater Plan do not provide any specific protection of piped stream habitats, the biota that inhabit them, or formally recognise them as migration pathways for freshwater fish. For example, if a section of piped stream was to be disturbed or replaced, there are no planning rules that would trigger the consideration of any ecological values or even alert GWRC freshwater scientists that the work was planned, even if the pipe in question is a fish migration pathway or has a permanent fish community within it. It would be sensible to work towards having piped stream recognised for their ecological values and be managed as such in the next iteration of the Natural Resources Plan.




Figure 25 A change in circular, concrete pipe dimensions in Pae Kawakawa Stream under The Parade in Island Bay in a high velocity section of piped stream has created a small, undercut drop that may impede fish passage.

5 RECOMMENDATIONS

5.1 Sampling Methodologies

- » Develop a rapid piped stream ecological assessment protocol that can be done from manholes without the need for a confined space entry. This would allow multiple sites to be covered in a day and sampling of pipes too small and/or too hazardous for entry. This could include:
 - Sampling of macroinvertebrates using a modified, long-handled kick net and multiple manhole entry points (more entry points will be needed to obtain sufficient representative samples).
 - Habitat assessment of visible section of piped stream including measurement or visual estimation of water depth, wetted width, water velocity, substrate composition, and organic matter type and abundance. This can be improved by lowering of a GoPro camera or similar on a pole to obtain footage of general habitat upstream and downstream of the manhole.
 - Observation of any eels that happen to be present and visible down manhole shafts.
- » Undertake detailed surveys that require pipe entry at key locations in main catchments. In practice this would likely only be in the middle to lower parts of larger catchments due to pipe size constraints. Any future detailed surveys should increase the area sampled for macroinvertebrates (e.g., collect pooled Surber samples rather than single samples) and include a fish survey via spotlighting and hand netting of at least 200 m of channel (so that actual fish species can be determined).

5.2 Habitat Improvements

- » Investigate the viability of undertaking various in-pipe habitat improvement retrofits including:
 - Installation of flexible baffles (or similar) in zones of high velocity, laminar flows to slow water velocities, increase water depths, and provide resting areas for resident and migrating fish.
 - Installation of artificial elements such as small pipes or half pipes securely attached to the pipe base to provide fish cover for resident and migrating fish.
 - Installation of structures to increase water depths, create pool habitat, and increase retention of organic and stony substrate material. These could take the form of leaky weirs constructed of small boulders at appropriate locations to create pools and small wood or rock vanes across the low flow channel.
- » Some locations have *in situ* materials from which fish cover can be constructed, however such improvised features are unlikely to withstand high flow events (Figure 26).
- » Where old pipes are being replaced or new piped sections constructed, investigate integrating in-pipe habitat features into the design rather than just using the standard, smooth bottomed concrete pipe design. This could include constructed pools, and designed low flow channels that incorporate fish cover elements and an irregular, rough pipe bottom to facilitate retention of stony bed substrates and organic matter.



Figure 26 Improvised fish cover feature constructed from rubble found at one piped stream site.

5.3 Recognition and Protection

- » Undertake a formal programme to identify sections of the stormwater pipe network that can be considered to be “piped streams” in that they are permanently flowing and close to where a natural, open stream was known to be or highly likely to have been located historically. This would involve extensive examination of WWL stormwater pipe GIS layers, waterway GIS layers, historical information on streams and urban development, and institutional knowledge at WWL, local councils, and the contractors who regularly work with the stormwater pipe network. The ultimate goal would be to create a detailed and accurate “piped stream” GIS layer.
- » After an accurate “piped stream” GIS layer has been created, initiate work to populate it with relevant metadata including presence or absence of permanent fish population, importance as a migratory fish pathway, in-pipe habitat quality, and macroinvertebrate community metrics. This would require a mix of GIS analyses and field investigations.
- » Aim to include provisions around maintaining and improving piped stream ecological values in the next iteration of the regional plan.

5.4 Further Research

- » A more widespread general survey of piped streams in the region, which should include smaller diameter pipes that would be unsuitable for human entry. This would feed into the recognition and protection recommendations above.
- » Initiate research into the length of piping that results in permanent changes to fish and macroinvertebrate assemblages. Unpublished data held by EOS Ecology from a 220 m pipe in Wellington indicates a macroinvertebrate and fish community in the pipe largely similar to that of natural open stream habitat upstream (i.e., abundant EPT taxa and a relatively diverse fish fauna (eels, banded kokopu, kōaro) resident in the pipe). There could be some pipe length threshold at which there is a shift towards only those fauna that can cope with piped stream conditions, such as those found in this pilot study. Because of its topography, the greater Wellington region is a good candidate to find sampleable pipes of different lengths with open channels upstream or downstream.
- » Initiate research into the importance of remnant open stream channels at maintaining catchment biodiversity, including the length of open habitat required for a macroinvertebrate and fish assemblage comprised of open stream habitat preferring taxa to be self-sustaining. This could help inform the minimum length of pipe daylighting required to attract open habitat preferring taxa. Part of such a study could involve determining how far downstream open habitat preferring taxa can persist in the pipe network and a comparison with longitudinal studies of natural cave systems (e.g., Watson, 2010).
- » A test of the effectiveness of in-pipe habitat improvements at suitable locations. This would include the design and installation of baffles at some locations and structures to increase water depths at others. Before-after, control-impact sampling of fish and macroinvertebrates could be used to determine if such features are worthwhile to install in piped streams.

6 MEDIA

- » Ecological work in piped streams has the ability to capture the imagination of journalists as a press release by GWRC prior to the fieldwork resulted in stories by TVNZ's One News and Radio New Zealand:
 - www.tvnz.co.nz/one-news/new-zealand/whitebait-eels-found-in-wellingtons-stormwater-system
 - www.rnz.co.nz/national/programmes/ourchangingworld/audio/2018697287/the-streams-beneath-the-streets

7 ACKNOWLEDGEMENTS

I greatly appreciated the assistance of various WWL staff for providing early guidance and GIS information (Michelle Chew, Rhyann McCoy, Paul Davison, Jack Furnish). Megan Oliver and Evan Harrison from GWRC provided much encouragement and funded the work, with assistance from Wellington City Council. Evan also provided helpful peer review comments. Michael Greer (formerly GWRC) helped get the project off the ground in the early stages. Thanks to Renee Wilkie and the crew from Silver Linings Contracting who ensured the fieldwork was safe, efficient, and enjoyable. Thank you to my EOS Ecology colleagues for peer review (Shelley McMurtrie), GIS assistance (Kirsty Brennan), and proof reading (Ariana Painter).

8 REFERENCES

- Clapcott, J. & Goodwin, E. 2014. Technical report of Macroinvertebrate Community Index predictions for the Greater Wellington region. Prepared for Greater Wellington Regional Council. Cawthron Report No. 2503. 20 p. plus appendices.
- Clarke, K.R. & Gorley, R.N. 2006. PRIMER v6 user manual/tutorial. PRIMER-E, Plymouth, UK. 190 p.
- James, A. 2016. Lambton Harbour catchment ICMP Stage 2: ecological assessment. EOS Ecology Report No. WEL02-15026-01. EOS Ecology, Christchurch, New Zealand. 59 p.
- James, A. 2017a. Evans Bay catchment ICMP Stage 2: ecological assessment. EOS Ecology Report No. WEL02-16073-01. EOS Ecology, Christchurch, New Zealand. 58 p.
- James, A. 2017b. South-east Coast catchment ICMP Stage 2: ecological assessment. EOS Ecology Report No. WEL02-16073-02. EOS Ecology, Christchurch, New Zealand. 53 p.
- James, A. 2018a. Island Bay-Houghton Bay catchment ICMP Stage 2: ecological assessment. EOS Ecology Report No. WEL02-16074-01. EOS Ecology, Christchurch, New Zealand. 51 p.
- James, A. 2018b. Lyall Bay catchment ICMP Stage 2: ecological assessment. EOS Ecology Report No. WEL02-16074-02. EOS Ecology, Christchurch, New Zealand. 40 p.
- James, A. 2019. Freshwater ecology of piped streams in Wellington: pilot study interim report. EOS Ecology Report No. GRE01-17087-01. 16 p.
- Harding, J.S., Clapcott, J.E., Quinn, J.M., Hayes, J.W., Joy, M.K., Storey, R.G., Greig, H.S., Hay, J., James, T., Beech, M.A., Ozane, R., Meredith, A.S. & Boothroyd, I. 2009. *Stream Habitat Assessment Protocols for Wadeable Rivers and Streams of New Zealand*. School of Biological Sciences, University of Canterbury, Christchurch, New Zealand. 133 p.
- Neale, M.W. & Moffett, E.R. 2016. Re-engineering buried urban streams: daylighting results in rapid changes in stream invertebrate communities. *Ecological Engineering* 87: 175-184.
- Stark, J.D. 1985. A macroinvertebrate community index of water quality for stony streams. Taranaki Catchment Commission, Wellington. Water & Soil Miscellaneous Publication No. 87. 53 p.
- Stark, J.D. & Maxted, J.R. 2007. A User Guide for the Macroinvertebrate Community Index. Cawthron Institute, Nelson. Report No. 1166. 66 p.
- Watson, T. 2010. The structuring of aquatic macroinvertebrate communities within cave streams. MSc thesis. University of Canterbury, Christchurch, New Zealand. 111 p.

9 APPENDICES

9.1 Appendix 1 – SIMPER Results Comparing Sites

Appendix Table 1 SIMPER results showing those taxa that contribute >5% of the differences between stream sites. Abundances shown are forth root transformed values. Also shown in parentheses are ANOSIM R-values for each pairwise comparison.

Taxon	Average Abundance	Average Abundance	% Contribution
The Parade–Dover St & Waipapa Stream (R=0.64, p=0.001)	The Parade–Dover St	Waipapa Stream	
Orthoclaadiinae midge larvae	1.26	0.19	18.02
Psychodidae gnat larvae	1.25	0.31	16.65
Oligochaeta worms	1.43	0.53	15.39
Collembola springtails	0.73	1.13	12.66
Scirtidae beetle larvae	0.58	0	8.82
<i>Polypedilum</i> midge larvae	0.43	0.14	6.31
Acarina mites	0.33	0.14	5.01
The Parade–Dover St & 302 The Parade (R=0.82, p=0.001)	The Parade–Dover St	302 The Parade	
Psychodidae gnat larvae	1.25	0	15.73
Orthoclaadiinae midge larvae	1.26	0	15.3
Oligochaeta worms	1.43	0.52	13.05
<i>Potamopyrgus</i> snails	0.17	1.16	12.43
<i>Physa</i> snails	0	0.8	8.53
Collembola springtails	0.73	0.09	8.38
Scirtidae beetle larvae	0.58	0	6.82
Waipapa Stream & 302 The Parade (R=0.69, p=0.001)	Waipapa Stream	302 The Parade	
Collembola springtails	1.13	0.09	23
<i>Potamopyrgus</i> snails	0	1.16	21.87
<i>Physa</i> snails	0.14	0.8	13.36
Oligochaeta worms	0.53	0.52	13.11
Psychodidae gnat larvae	0.31	0	5.1
<i>Ferrissia</i> limpets	0	0.29	5.08
The Parade–Dover St & 348 The Parade (R=0.93, p=0.001)	The Parade–Dover St	348 The Parade	
<i>Potamopyrgus</i> snails	0.17	2.66	21.48
Psychodidae gnat larvae	1.25	0	11.23
Orthoclaadiinae midge larvae	1.26	0.16	10.66
<i>Ferrissia</i> limpets	0	0.95	8.91
Oligochaeta worms	1.43	1.46	8.75
<i>Physa</i> snails	0	1.05	8.48
Collembola springtails	0.73	0.1	6.18
Scirtidae beetle larvae	0.58	0	5.02
Waipapa Stream & 348 The Parade (R=0.92, p=0.001)	Waipapa Stream	348 The Parade	

<i>Potamopyrgus</i> snails	0	2.66	28.11
Oligochaeta worms	0.53	1.46	12.99
Collembola springtails	1.13	0.1	12.71
Acarina mites	0	0.95	11.79
<i>Physa</i> snails	0.14	1.05	9.91
302 The Parade & 348 The Parade (R=0.27, p=0.002)	302 The Parade	348 The Parade	
<i>Potamopyrgus</i> snails	1.16	2.66	27.21
Oligochaeta worms	0.52	1.46	18.6
<i>Ferrissia</i> limpets	0.29	0.95	14.31
<i>Physa</i> snails	0.8	1.05	12.88
Acarina mites	0.11	0.3	5.86
The Parade–Dover St & Miramar-Shops (R=0.68, p=0.001)	The Parade–Dover St	Miramar-Shops	
Psychodidae gnat larvae	1.25	0	16.99
Orthoclaadiinae midge larvae	1.26	0.27	13.32
<i>Polypedilum</i> midge larvae	0.43	1.07	11.85
Collembola springtails	0.73	0	9.16
Oligochaeta worms	1.43	1.42	8.56
Acarina mites	0.33	0.79	8.54
Scirtidae beetle larvae	0.58	0	7.51
<i>Potamopyrgus</i> snails	0.17	0.53	6.68
Waipapa Stream & Miramar-Shops (R=0.74, p=0.001)	Waipapa Stream	Miramar-Shops	
Collembola springtails	1.13	0	19.22
Oligochaeta worms	0.53	1.42	17.34
<i>Polypedilum</i> midge larvae	0.14	1.07	15.58
Acarina mites	0.14	0.79	11.03
<i>Potamopyrgus</i> snails	0	0.53	7.32
Platyhelminthes flatworms	0	0.29	5.61
Orthoclaadiinae midge larvae	0.19	0.27	5.17
302 The Parade & Miramar-Shops (R=0.51, p=0.001)	302 The Parade	Miramar-Shops	
Oligochaeta worms	0.52	1.42	19.04
<i>Polypedilum</i> midge larvae	0.09	1.07	16.64
<i>Potamopyrgus</i> snails	1.16	0.53	14.7
Acarina mites	0.11	0.79	11.86
<i>Physa</i> snails	0.8	0	11.45
Platyhelminthes flatworms	0	0.29	5.89
348 The Parade & Miramar-Shops (R=0.70, p=0.001)	348 The Parade	Miramar-Shops	
<i>Potamopyrgus</i> snails	2.66	0.53	24.3
<i>Polypedilum</i> midge larvae	0	1.07	11.57
Oligochaeta worms	1.46	1.42	11.54
<i>Ferrissia</i> limpets	0.95	0	11.47
<i>Physa</i> snails	1.05	0	10.62
Acarina mites	0.3	0.79	7.71

The Parade–Dover St & Miramar Park (R=0.98, p=0.001)	The Parade–Dover St	Miramar Park	
<i>Potamopyrgus</i> snails	0.17	1.41	15.74
Psychodidae gnat larvae	1.25	0	15.63
Orthoclaadiinae midge larvae	1.26	0	15.28
<i>Physa</i> snails	0	0.73	9.03
Collembola springtails	0.73	0.08	8.45
Scirtidae beetle larvae	0.58	0	6.9
Oligochaeta worms	1.43	1.57	6.65
Waipapa Stream & Miramar Park (R=0.94, p=0.001)	Waipapa Stream	Miramar Park	
<i>Potamopyrgus</i> snails	0	1.41	23.19
Oligochaeta worms	0.53	1.57	19.67
Collembola springtails	1.13	0.08	18.73
<i>Physa</i> snails	0.14	0.73	11.64
302 The Parade & Miramar Park (R=0.20, p=0.004)	302 The Parade	Miramar Park	
Oligochaeta worms	0.52	1.57	31.38
<i>Physa</i> snails	0.8	0.73	18.9
<i>Potamopyrgus</i> snails	1.16	1.41	18.26
<i>Ferrissia</i> limpets	0.29	0.28	10.27
348 The Parade & Miramar Park (R=0.35, p=0.001)	348 The Parade	Miramar Park	
<i>Potamopyrgus</i> snails	2.66	1.41	25.42
Oligochaeta worms	1.46	1.57	16.77
<i>Ferrissia</i> limpets	0.95	0.28	14.99
<i>Physa</i> snails	1.05	0.73	12.01
Sphaeriidae pea clams	0.35	0.08	5.44
Acarina mites	0.3	0	5.37
Miramar-Shops & Miramar Park (R=0.60, p=0.001)	Miramar-Shops	Miramar Park	
<i>Potamopyrgus</i> snails	0.53	1.41	17.9
<i>Polypedilum</i> midge larvae	1.07	0.08	17.18
<i>Physa</i> snails	0	0.73	12.79
Acarina mites	0.79	0	12.66
Oligochaeta worms	1.42	1.57	9.7
Platyhelminthes flatworms	0.29	0	5.73

9.2 Appendix 2 - BEST (Biota and Environment Matching) Results

Appendix Table 2 BEST results from piped stream sites showing those measured habitat parameters that best matched the observed macroinvertebrate community composition. Habitat variables included in the analysis were water depth, wetted width, water velocity, biofilm cover, and substrate index.

Variables	Correlation
Wetted width, Water velocity	0.356
Water velocity	0.320
Wetted width, Water velocity, Biofilm cover	0.312
Wetted width, Water velocity, Substrate index	0.287
Water velocity, Biofilm cover	0.282
Wetted width, Water velocity, Biofilm cover, Substrate index	0.281
Water velocity, Substrate index	0.271
Water velocity, Biofilm cover, Substrate index	0.256
Water depth, Wetted width, Water velocity	0.248
Water depth, Water velocity	0.233

9.3 Appendix 3 – SIMPER Results Comparing Catchments

Appendix Table 3 SIMPER results showing those taxa that contribute >5% of the differences between catchments (Pae Kawakawa, Waipapa, and Miramar). Abundances shown are forth root transformed values. Also shown in parentheses are ANOSIM R-values and significance level for each pairwise comparison.

Taxon	Average Abundance	Average Abundance	% Contribution
Pae Kawakawa & Waipapa (R=0.35, p=0.001)	Pae Kawakawa	Waipapa Stream	
<i>Potamopyrgus</i> snails	1.33	0	18.54
Collembola springtails	0.31	1.13	16.19
Oligochaeta worms	1.15	0.53	13.72
<i>Physa</i> snails	0.61	0.14	8.87
Psychodidae gnat larvae	0.43	0.31	7.67
Orthoclaadiinae midge larvae	0.49	0.19	7.21
<i>Ferrissia</i> limpets	0.42	0	6.13
Pae Kawakawa & Miramar (R=0.05, p=0.096)	Pae Kawakawa	Miramar	
<i>Potamopyrgus</i> snails	1.33	0.99	17.29
Oligochaeta worms	1.15	1.5	14.74
<i>Physa</i> snails	0.61	0.38	9.96
<i>Polypedilum</i> midge larvae	0.18	0.56	8.68
Orthoclaadiinae midge	0.49	0.13	7.17
<i>Ferrissia</i> limpets	0.42	0.14	6.98
Acarina mites	0.25	0.38	6.62
Psychodidae gnat larvae	0.43	0	5.96
Waipapa & Miramar (R=0.73, p=0.001)	Waipapa	Miramar	
Collembola springtails	1.13	0.04	18.96
Oligochaeta worms	0.53	1.5	18.55
<i>Potamopyrgus</i> snails	0	0.99	15.54
<i>Polypedilum</i> midge larvae	0.14	0.56	9.2
<i>Physa</i> snails	0.14	0.38	7.1
Acarina mites	0.14	0.38	6.2

9.4 Appendix 4 – SIMPER Results Comparing Open and Piped Stream Sites

Appendix Table 4 SIMPER results showing those taxa that contribute >5% of the differences between open and piped stream sites in the Pae Kawakawa and Miramar catchments. Abundances shown are relative abundances. Also shown in parentheses are ANOSIM R-values and significance level.

Taxon	Average Abundance	Average Abundance	% Contribution
Open & Piped (R=0.56, p=0.048)	Open	Piped	
Oligochaeta worms	4.59	33.92	20.04
<i>Potamopyrgus</i> snails	21.27	33.3	18.38
<i>Polypedilum</i> midge larvae	14.98	7.22	10.05
<i>Hydropsyche</i> (<i>Orthopsyche</i>) caddisfly larvae	12.42	0.01	8.48
Orthoclaadiinae midge larvae	13.52	3.38	8.39
<i>Physa</i> snails	0.02	7.65	5.23

www.eosecology.co.nz
info@eosecology.co.nz

PO Box 4262, Christchurch 8140
P 03 389 0538

PO Box 8054, Palmerston North 4446
P 06 358 9566

SCIENCE + ENGAGEMENT

